


Rasbawati Rasbawati

Impact of Golden Banana Puree (*Musa acuminata*) Enrichment on the pH, Lactic Acid Bacteria Count, and Viscosity of Kefir

 JURNAL DOSEN

 LP2M

 Universitas Muhammadiyah Pare-Pare

Document Details

Submission ID

trn:oid:::1:3260270135

Submission Date

May 26, 2025, 9:07 AM GMT+7

Download Date

May 26, 2025, 9:12 AM GMT+7

File Name

Agripet_2024.pdf

File Size

196.7 KB

7 Pages

4,663 Words

25,086 Characters

14% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.





Filtered from the Report

- Bibliography
- Small Matches (less than 9 words)




Exclusions

- 16 Excluded Sources

Match Groups

-  **23 Not Cited or Quoted 12%**
Matches with neither in-text citation nor quotation marks
-  **4 Missing Quotations 2%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 12%  Internet sources
- 8%  Publications
- 5%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 23 Not Cited or Quoted 12%**
Matches with neither in-text citation nor quotation marks
- 4 Missing Quotations 2%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 12% Internet sources
- 8% Publications
- 5% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	jitek.ub.ac.id	1%
2	Student papers	Syiah Kuala University	1%
3	Internet	pmc.ncbi.nlm.nih.gov	1%
4	Internet	jnp.fapet.unsoed.ac.id	<1%
5	Internet	ojs.unud.ac.id	<1%
6	Publication	Helmi Musyaffa Akmal, Juni Sumarmono, Triana Setyawardani. "Pengaruh Penam...	<1%
7	Internet	fsct.modares.ac.ir	<1%
8	Internet	pjlss.edu.pk	<1%
9	Publication	Lusia Yotista Enggal Parasthi, Diana Nur Afifah, Choirun Nissa, Binar Panunggal. ...	<1%
10	Internet	www.ncbi.nlm.nih.gov	<1%

11	Student papers	Universitas Sebelas Maret	<1%
12	Internet	ojs.umb-bungo.ac.id	<1%
13	Internet	biotecnia.unison.mx	<1%
14	Internet	sbp-journal.com	<1%
15	Publication	Dinda Sukma Wibowo, Maria Marina Herawati. "The Effect of Application of Low-c...	<1%
16	Internet	download.bibis.ir	<1%
17	Internet	www.mdpi.com	<1%
18	Internet	journal.ugm.ac.id	<1%
19	Internet	jurnalfkip.unram.ac.id	<1%
20	Internet	ejournal2.undip.ac.id	<1%
21	Internet	jap.fapet.unsoed.ac.id	<1%
22	Internet	perpustakaan.poltekkes-malang.ac.id	<1%
23	Publication	"Microbes in the Food Industry", Wiley, 2023	<1%
24	Publication	M. Fiqri Erzhad, Reza Adiyoga, Marwah H, Zakiah Wulandari, M Sriduresta Soenar...	<1%

25

Internet

ir.niist.res.in:8080

<1%



Impact of Golden Banana Puree (*Musa acuminata*) Enrichment on the pH, Lactic Acid Bacteria Count, and Viscosity of Kefir

Rasbawati^{1*}, Fitriani², Oktavia Rahayu Puspitarini³, Intan Dwi Novieta¹, Irmayani¹

¹Department of Animal Husbandry, Faculty of Agriculture, Animal Husbandry and Fisheries, Muhammadiyah University Parepare, Indonesia

²Department of Animal Husbandry, Pangkajene Islands State Agricultural Polytechnic, Indonesia

³Faculty of Animal Husbandry, Islamic University of Malang, Indonesia

ABSTRACT. The addition of golden banana puree (*Musa acuminata*) as a prebiotic has been hypothesized to enhance the growth of lactic acid bacteria in kefir. This study aimed to determine the effects of adding golden banana puree on the pH value, total lactic acid bacteria (LAB) count, and viscosity of kefir. The experiment was conducted using a Completely Randomized Design (CRD) with four treatments and three replications. The treatments included the addition of 0% (P0), 6% (P1), 8% (P2), and 10% (P3) banana puree. The observed variables were pH value, total LAB, and viscosity. Data were analyzed using Analysis of Variance (ANOVA), followed by Duncan's Multiple Range Test (DMRT) when a significant treatment effect was detected. The results indicated that the addition of golden banana puree had a very significant effect ($P < 0.01$) on the pH value, total LAB count, and viscosity of kefir. The study concluded that increasing the concentration of golden banana puree leads to a higher total LAB count and viscosity, while inversely lowering the pH value.

Keywords: golden banana puree, viscosity, kefir, pH value, total LAB

INTRODUCTION

In recent years, particularly following the Covid-19 pandemic, there has been an increased public awareness of health and nutrition. This has led to a growing demand for products that offer not only nutritional benefits but also positive health effects, commonly referred to as functional foods (Martharini and Indratiningsih, 2017). Kefir, a fermented drink with a distinctive sour and slightly alcoholic taste and a creamy consistency, is one such functional food (Yousefvand et al., 2022). Originating from the Caucasus and Tibet (Rosa et al., 2017), kefir is produced by fermenting milk (cow, goat, or sheep) with a mix of lactic acid bacteria (LAB), acetic acid bacteria, and yeast present in kefir grains (Ningsih et al., 2018). Kefir is known for its probiotic properties and is easily digestible (Łopusiewicz et al., 2022).

Kefir grains contain a complex community of microorganisms, including LAB such as *Lactobacillus* spp., *Leuconostoc* spp., *Streptococcus* spp., and *Lactococcus* spp., as well as yeasts like *Saccharomyces* spp., *Kluyveromyces* spp., *Zygosaccharomyces* spp., *Torulopsis* spp., and *Candida* spp., and possibly acetic acid bacteria (*Acetobacter* spp.) (Rohmah and Estiasih, 2018).

Kefir is classified as a functional food due to its bioactive components, which offer various health benefits. These include the presence of beneficial bacteria, vitamins, minerals, and

essential amino acids that contribute to overall health and well-being (Kartika et al., 2020; Julianto et al., 2016). Research has shown that kefir can have positive effects on various health conditions, including colorectal cancer, cardiovascular disease, type 2 diabetes mellitus, obesity, and kidney disease (Yilmaz et al., 2022). Additional benefits include reducing inflammatory bowel disease severity (Yilmaz et al., 2019), exhibiting anticarcinogenic properties (Sharifi et al., 2017), improving serum lipid profiles (Fathi et al., 2017), and mitigating neurodegenerative diseases (Ton et al., 2020). Kefir is also used in cosmetic applications for skin lightening, anti-acne treatments, wound healing, and as an antioxidant (Dewi et al., 2018).

Despite its numerous benefits, kefir's consumption is often limited by its sour taste. To enhance its appeal, it is beneficial to explore the incorporation of other ingredients. Golden banana (*Musa acuminata*), which is rich in nutrients and acts as a prebiotic, could be an ideal candidate for this purpose. Golden banana, a monocot plant thriving in tropical regions such as Indonesia (Puguh et al., 2020), has high production yields, contributing significantly to the fruit market in Indonesia (Junior and Hariyono, 2020). Known for its round shape, bright yellow skin, and sweet flavor, golden banana offers several health benefits. According to the Indonesian Food Composition Table (TKPI) (2019), it contains essential nutrients including carbohydrates, fiber, vitamins, and minerals. It is particularly noted for its potential to reduce LDL cholesterol and increase HDL cholesterol, thereby improving cardiovascular


*Corresponding author: rasbawati@gmail.com

Received: 18 August 2022

Revised: 15 January 2024

Accepted: 31 July 2024

DOI: <https://doi.org/10.17969/agripet.v24i2.27562>



health and reducing hypertension risk (Badriah *et al.*, 2019). However, challenges related to its short shelf life and handling necessitate research into incorporating golden banana puree into kefir production.

This study aims to evaluate the effects of adding golden banana (*Musa acuminata*) puree on the pH value, total lactic acid bacteria (LAB) count, and viscosity of kefir.

MATERIALS AND METHODS

Material

The materials used in this study included pure milk sourced from dairy farms in Enrekang Regency, commercially obtained kefir grains, golden bananas, MRSA, distilled water, 70% alcohol, and other laboratory supplies such as tissue and label paper. The equipment consisted of a pan, stove, measuring cup, thermometer, digital scale, petri dish, containers, 200 and 400 mesh filters, pH meter, incubator, pycnometer, blender, and stirrer.

Method

The research employed an experimental approach using a Completely Randomized Design (CRD) with four treatments and three replications. The treatments were as follows: P0 (0% golden banana puree), P1 (6% golden banana puree), P2 (8% golden banana puree), and P3 (10% golden banana puree). The observed variables included pH value, total lactic acid bacteria (LAB) count, and viscosity.

Research Procedure

Development of Banana Puree

Golden banana puree was prepared following the methods of Kartika *et al.* (2020) and Tuhumury *et al.* (2018), with some modifications. Fully ripe golden bananas with yellow skin were sorted, cleaned, and steamed for 5 minutes before peeling. After steaming, the bananas were mashed using a blender to obtain the puree.

Development of Kefir

Kefir was produced by adding banana puree according to the procedure outlined by Mulyani *et al.* (2021) with modifications. One liter of milk was pasteurized for 30 minutes at 65°C. After cooling to room temperature, banana puree was added at the specified treatment levels (6%, 8%, and 10% of milk volume). The mixture was stirred gently, then inoculated with 5% kefir grains aseptically and stirred again. The milk was incubated at room temperature (20-25°C) for 24 hours. After

incubation, the kefir was filtered through a 40-mesh nylon sieve to separate the kefir grains (Purba *et al.*, 2018).

pH Value Measurement

The pH value was measured using a pH meter. The electrode was cleaned with distilled water and calibrated with pH buffer solutions of 4 and 7 before use. The electrode tip was then inserted into the kefir sample until the pH meter displayed a stable reading, which was recorded (Prastujati *et al.*, 2018).

Measurement of Total Lactic Acid Bacteria (LAB)

The total LAB count was determined using the Total Plate Count (TPC) method. Dilutions of 10⁻⁵ and 10⁻⁶ were prepared. One milliliter of each dilution was plated onto sterile deMan Rogosa Sharpe Agar (MRSA) and mixed using the pour plate method. The plates were incubated for 24 hours at 37°C. After incubation, colonies were counted (Sinurat *et al.*, 2018).

Viscosity Measurement

Viscosity was measured using the Ostwald pipette method as described by Prastiwi *et al.* (2018). Initially, the specific gravity of the milk was determined by weighing an empty pycnometer, adding 10 ml of the sample, and weighing it again. For viscosity measurement, 10 ml of the sample was placed into an Ostwald pipette. The time taken for the sample to descend from the top to the bottom mark was recorded using a stopwatch. The viscosity was then calculated using the formula:

$$\text{viscositas} = \frac{\rho \times t_{\text{sample}} \times \eta_{\text{water}}}{\rho \times t_{\text{water}}}$$

$$\rho_{\text{sample}} = \frac{m - m}{v}$$

information :

m : mass of the empty pycnometer (g)

m : mass of pycnometer + sample (g)

v : pycnometer volume (ml)

η : water viscosity

ρ : density (g/ml) (water = 1 g/ml)

t : time (s)

Data Analysis

Data were analyzed using Analysis of Variance (ANOVA) to determine the effects of different treatments on pH value, total lactic acid bacteria (LAB), and viscosity. When significant effects were detected, the Duncan Multiple Range

Test (DMRT) was performed to compare means between treatments. Statistical analysis was conducted using SPSS version 16.0 for Windows with a significance level set at 1%.

RESULTS AND DISCUSSION

The results for pH values, total lactic acid bacteria, and kefir viscosity with the addition of golden banana puree (*Musa acuminata*) at levels of 0%, 6%, 8%, and 10% are summarized in Table 1.

Table 1. pH value, total lactic acid bacteria and kefir viscosity the with addition banana puree

Variable	Treatment			
	P0	P1	P2	P3
pH	4.93±0.15 ^c	4.57±0.06 ^b	4.43±0.06 ^b	4.03±0.06 ^a
Total LAB (CFU/gr)	1.1x10 ⁴ ±0.207 ^a	8.9x10 ⁴ ±0.065 ^b	1.3x10 ⁵ ±0.049 ^b	2.7x10 ⁷ ±0.086 ^c
Viscosity (cp)	8.78±0.95 ^a	12.17±0.76 ^{ab}	13.63±0.68 ^b	14.93±3.81 ^b

Note: ^{a, b, c} different lowercase superscript letters indicate very significant differences (P<0.01)

pH Value

The results showed that the pH values for kefir with golden banana puree at different concentrations were as follows: 4.93 for the control (P0), 4.57 for 6% banana puree (P1), 4.43 for 8% banana puree (P2), and 4.03 for 10% banana puree (P3) (Table 1). The analysis of variance revealed that the addition of golden banana puree significantly affected the pH value of kefir (P<0.01). The kefir with 10% banana puree exhibited the lowest pH (4.03), while the control had the highest pH (4.93).

The significant difference in pH values indicates that kefir with 10% golden banana puree is more acidic compared to the control and other treatments. The control treatment had the highest pH value due to the absence of additional carbohydrates. Without extra carbohydrates, the fermentation media had limited nutrients for lactic acid bacteria (LAB) to produce lactic acid, leading to a higher pH (Rizqiati *et al.*, 2021).

As the concentration of banana puree increased, the pH value decreased. This is attributed to the carbohydrates present in bananas, which are fermented by LAB into lactic acid. The production of lactic acid during fermentation increases the acidity and lowers the pH value (Utomo *et al.*, 2019; Rasbawati *et al.*, 2019). Lactose, the primary carbohydrate in milk, is also broken down into lactic acid during fermentation, contributing to the lower pH value as the fermentation progresses (Martharini and Indratiningsih, 2017).

The trend observed in this study aligns with findings from Barus *et al.* (2019), who noted that higher lactic acid production results in a more sour taste and lower pH. The pH values observed in this study are consistent with those reported by Yousefvand *et al.* (2022), where kefir pH on the first day of fermentation ranged between 4.49 and

4.53. Similarly, Kartika *et al.* (2020) found that increasing concentrations of gembili fruit puree led to a decrease in pH values. The pH of fermented beverages is influenced by organic acids such as lactic acid, acetic acid, butyric acid, and propionic acid produced during fermentation (Viogenta *et al.*, 2021).

Total LAB

Lactic acid bacteria are the main microbes used in milk fermentation (Savaiano and Hutkins, 2021). Data on LAB kefir with the addition of golden banana with different levels can be seen in Table 1. The addition of golden bananas at different levels had a very significant effect (P<0.01) on the total LAB of kefir. Table 1 shows that the lowest total LAB was in the control treatment (2.7x10⁷ CFU/g) and the highest total LAB was in the 10% golden banana treatment (2.7x10⁷ CFU/g). The higher the golden banana concentration (up to 10%), the greater the LAB growth. This phenomenon occurs allegedly due to the addition of golden bananas that contains glucose and fructose was able to support the development of LAB. In other words, the addition of golden bananas can increase the total activity of lactic acid bacteria so that the number of microbes in kefir increases. According to Meghwar *et al.* (2021) bananas contain fructose where it was used as an energy source for the LAB starter so that the growth was optimal during the fermentation process and was able to breaks down lactose into lactic acid. This is in accordance with the statement by (Pratangga, D *et al.*, 2019) that LAB utilizes the sugar in the fermentation media for its growth. The utilization of sugars in the substrate for LAB growth can be seen by increasing the LAB cell population. Sucrose from sugar will be broken down by yeast using the enzyme invertase to produce glucose and fructose, then the results will be utilized by lactic acid bacteria and yeast to produce pyruvic acid

through the glycolysis process. The pyruvic acid obtained from the glycolysis process will be converted into lactic acid and will be converted into alcohol and carbon dioxide (Mulyani *et al.*, 2021). Sucrose in bananas and lactose in milk are part of the disaccharides which play an important role for LAB to carry out the fermentation process. LAB growth was influenced by several factors including free water content (a_w), oxygen, availability of substrate in the growth medium, environmental temperature and the initial pathogenic microbes (Pratangga, D *et al.*, 2019).

The results of this study was similar to (Agustine *et al.*, 2018) that yoghurt with additional levels of sucrose and skim milk had increased results, this indicates that the higher variation of sucrose and skim milk will resulting in higher LAB activity due to the presence of sucrose which is a carbon source for the growth of microorganisms.

Viscosity

Lactic acid bacteria (LAB) are crucial for the fermentation of milk, as they produce lactic acid that affects the texture and flavor of kefir (Savaiano and Hutkins, 2021). The data on LAB counts in kefir with varying levels of golden banana puree are presented in Table 1. The addition of golden banana puree significantly influenced the total LAB count ($P < 0.01$). The control treatment (P0) had the lowest LAB count (2.7×10^7 CFU/g), while the kefir with 10% golden banana puree (P3) had the highest LAB count (2.7×10^7 CFU/g). The increase in LAB count with higher banana puree concentrations can be attributed to the presence of glucose and fructose in bananas, which support LAB growth.

Bananas provide fructose, which serves as an energy source for LAB, thereby enhancing their growth during fermentation (Meghwar *et al.*, 2021). LAB utilize the sugars present in the fermentation medium for growth, which is evident from the increase in LAB counts. Sucrose in bananas and lactose in milk are disaccharides that are broken down during fermentation, aiding in LAB growth and lactic acid production (Mulyani *et al.*, 2021). The presence of sucrose and lactose as carbon sources plays a crucial role in LAB fermentation processes (Pratangga, D *et al.*, 2019). This is similar to findings by Agustine *et al.* (2018), who observed that increasing sucrose and skim milk levels in yogurt led to higher LAB activity.

The viscosity of kefir increased significantly with the addition of golden banana puree ($P < 0.01$). The control treatment (P0) had the lowest viscosity (8.78 cP), while the 10% banana puree treatment

(P3) had the highest viscosity (14.93 cP). This increase in viscosity with higher banana puree concentrations can be attributed to the protein content in milk and the effects of banana puree. Proteins in milk bind water, which increases viscosity (Syafitri and Swarastuti, 2013). The presence of golden banana puree enhances the thickening effect as the proteins in milk coagulate due to lactic acid produced during fermentation. This coagulation and thickening are due to proteins reaching their isoelectric point at a pH of 4.7.

Similar findings were reported by Nurubay *et al.* (2021), who observed increased viscosity in goat's milk kefir with added bile banana puree. Paredes *et al.* (2022) also found increased viscosity in kefir with the addition of fruit and vegetable juices. Kefir grains, which contain LAB and yeast, contribute to viscosity through the production of exopolysaccharides (Jurášková *et al.*, 2022). Exopolysaccharides, produced by LAB, enhance the texture and flavor of kefir and provide several health benefits, including wound healing and immunomodulatory effects (Angelin and Kavitha, 2020; Hussain *et al.*, 2017). LAB strains such as *Lactococcus*, *Bifidobacterium*, and others are known for producing exopolysaccharides (Sanalibaba and Çakmak, 2016). Additionally, factors such as temperature, incubation time, and the concentration of raw materials affect the viscosity of kefir (Putri *et al.*, 2020).

CONCLUSION

The study concluded that increasing the concentration of golden banana (*Musa acuminata*) puree in kefir results in higher total lactic acid bacteria (LAB) counts and viscosity, while simultaneously decreasing the pH value.

ACKNOWLEDGEMENT

The author extends gratitude to Muhammadiyah University of Parepare for funding this research under the novice lecturer research scheme for the 2022 fiscal year.

REFERENCE

- Agustine, L., Okfrianti, Y., Jum, J., 2018. Identifikasi Total Bakteri Asam Laktat (BAL) pada Yoghurt dengan Variasi Sukrosa dan Susu Skim. Jurnal Dunia Gizi 1.
- Angelin J, Kavitha M. Exopolysaccharides from probiotic bacteria and their health potential. Int J Biol Macromol. 2020 Nov 1; 162:853-

865. doi: 10.1016/j.ijbiomac.2020.06.190. Epub 2020 Jun 22. PMID: 32585269; PMCID: PMC7308007.
- Badriah, D. L., Wulan, N., Yulianti, A., 2019. Pengaruh Konsumsi Pisang Emas (*Musa Acuminata*) Terhadap Penurunan Tekanan Darah Pada Klien Hipertensi Ringan di Kecamatan Selajambe Kabupaten Kuningan Tahun 2018. *Jurnal Ilmu Kesehatan Bhakti Husada: Health Sciences Journal*, 10(1), 11–17. <https://doi.org/10.34305/jikbh.v10i1.78>
- Barus, E.P.B., Rizqiaty, H., Bintoro, V.P., 2019. Total bakteri asam laktat, nilai pH, total padatan terlarut, dan sifat organoleptik cocolir dengan lama fermentasi yang berbeda. *J. Teknologi Pangan*, 3(2); 247–252
- Czarina, M., Moreno, R., Emata, O. C., 2020. Chemical and microbiological properties of kefir produced by kefir grains in raw and pasteurized cow's milk. *Mindanao Journal of Science and Technology*, 18(2), 277–292
- Dewi, M. L., Rusdiana, T., Muchtaridi, M., Putriana, N. A., 2018. Artikel Tinjauan: Manfaat Kefir untuk Kesehatan Kulit. *Farmaka*, 16(2), 80–86. <https://doi.org/10.24198/jf.v16i2.18052.g8478>
- Hussain, A., Zia, K.M., Tabasum, S., Noreen, A., Ali, M., Iqbal, R., Zuber, M., 2017. Blends and composites of exopolysaccharides; properties and applications: A review. *Int J Biol Macromol* 94, 10–27.
- İ. Yılmaz, M. E. Dolar, and H. Özpinar., 2019. Effect of administering kefir on the changes in fecal microbiota and symptoms of inflammatory bowel disease: a randomized controlled trial, *Turkish Journal of Gastroenterology*, vol. 30, no. 3, pp. 242–253
- Julianto, B., Rossi, E., Yusmarini, Y., 2016. Karakteristik Kimiawi dan Mikrobiologi Kefir Susu Sapi dengan Penambahan Susu Kedelai. *Jurnal Online Mahasiswa (JOM) Bidang Pertanian*, 3(1), 1–11. <https://jom.unri.ac.id/index.php/JOMFA-PERTA/article/view/9588>
- Junior, D. C., Hariyono, D., 2020. Kajian Hubungan Curah Hujan Dengan Produktivitas Tanaman Pisang Mas Kirana (*Musa Acuminata*) di Kabupaten Lumajang. *Jurnal Produksi Tanaman*, 8(2), 256–263. Retrieved from <http://protan.studentjournal.ub.ac.id/index.php/protan/article/view/1374>
- Jurášková, D., Ribeiro, S. C., Silva, C. C. G., 2022. January 1). Exopolysaccharides Produced by Lactic Acid Bacteria: From Biosynthesis to Health-Promoting Properties. *Foods*. MDPI. <https://doi.org/10.3390/foods11020156>
- Kartika, Rahayuningsih, M., Setyaningsih, D., 2020. Production and characterization of synbiotic kefir cow milk with addition of *Dioscorea esculenta* puree. In *IOP Conference Series: Earth and Environmental Science* (Vol. 472). Institute of Physics Publishing. <https://doi.org/10.1088/1755-1315/472/1/012026>
- Kemenkes RI., 2019. Tabel Komposisi Pangan Indonesia (TKPI) “Pisang mas segar. https://www.andrafarm.com/_andra.php?_i=0-home. Diunduh 10/08/2022
- Łopusiewicz, Ł., Śmietana, N., Paradowska, D., Drożdowska, E., 2022. Black Cumin (*Nigella sativa* L.) Seed Press Cake as a Novel Material for the Development of New Non-Dairy Beverage Fermented with Kefir Grains. *Microorganisms*, 10(2). <https://doi.org/10.3390/microorganisms10020300>
- M. Sharifi, A. Moridnia, D. Mortazavi, M. Salehi, M. Bagheri, and A. Sheikhi., 2017. Kefir: a powerful probiotics with anticancer properties, *Medical Oncology*, vol. 34, no. 11, p. 183
- Martharini, D., Indratiningsih, I., 2017. Kualitas Mikrobiologis dan Kimiawi Kefir Susu Kambing dengan Penambahan *Lactobacillus acidophilus* FNCC 0051 dan Tepung Kulit Pisang Kepok (*Musa Paradisiaca*). *agriTECH* 37, 23–30.
- Meghwar, P, A. B. Baloch, A. A. Depar, D. Kumar, S. M. Zahra, and R. Noreen., 2021. An Overview on Vital role of Banana and its Valorization. *International Journal of Food Chemistry and Human Nutrition Mini Review Article*.
- Mulyani, S., Sunarko, K.M.F., Setiani, B.E., 2021. Pengaruh Lama Fermentasi terhadap Total Asam, Total Bakteri Asam Laktat dan Warna Kefir Belimbing Manis (*Averrhoa carambola*). *Jurnal Ilmiah Sains* 21, 113.

- Mundiri, N. A., I. Megantara, T. T. K. Anggaeni., 2020. Kajian Pustaka: Pemanfaatan Eksopolisakarida Bakteri Asam Laktat Probiotik Asal Produk Pangan Fermentasi Sebagai Imunomodulator. Indonesia Medicus Veterinus. Pissn: 2301-7848; Eissn: 2477-6637. Doi: 10.19087/imv.2020.9.5.849
- Ningsih, D.R., Bintoro, V.P. Nurwantoro., 2018. Analisis total padatan terlarut, kadar alkohol, nilai pH dan total asam pada kefir optima dengan penambahan high fructose syrup (HFS). J. Teknologi Pangan, 2(2): 84-88.
- Nurubay. B. N, S. Saloko, dan M. D. Ariyana., 2021. Pengaruh Konsentrasi Puree Pisang Bile (*Musa parasidiaca*) Terhadap Sifat Mikrobiologi, Kimia dan Sensoris Kefir Susu Kambing. J. Teknologi Pertanian Andalas. Vol. 25 No. 2., September 2021, ISSN 1410-1920, EISSN 2579-4019
- P. Sanalibaba, G.A. Cakmak, Exopolysaccharides production by lactic acid bacteria, Appl. Microbiol. Open Access. 2., 2016. <https://doi.org/10.4172/2471-9315.1000115>
- Paredes, J. L., Escudero-Gilete, M. L., Vicario, I. M., 2022. A new functional kefir fermented beverage obtained from fruit and vegetable juice: Development and characterization. LWT, 154. <https://doi.org/10.1016/j.lwt.2021.112728>
- Prabowo, M. R. E., Sari, N. K., Zulaika, E., 2020. Kajian Yuridis Pendaftaran Pisang Mas Kirana Sebagai Produk Indikasi Geografis Kabupaten Lumajang. Syntax Idea, 2(9), 626–634
- Prastiwi, V. F., Bintoro, V. P., Rizqiati, H., 2018. Sifat Mikrobiologi, Nilai Viskositas dan Organoleptik Kefir Optima dengan Penambahan High Fructose Syrup (HFS). Jurnal Teknologi Pangan, 2(1), 27–32
- Prastujati, A. U., Hilmi, M., Khirzin, M. H., 2018. Pengaruh Konsentrasi Starter terhadap Kadar Alkohol, pH, dan Total Asam Tertitrasi (TAT) Whey Kefir. Jurnal Ilmu Peternakan Terapan, 1(2), 63–69.
- Pratangga, D. A., Susilowati, S., Puspitarini, O. R., 2019. Pengaruh Penambahan Berbagai Level Sukrosa dan Fruktosa terhadap Total Bakteri Asam Laktat dan Nilai pH Yoghurt Susu Kambing. Jurnal Rekasatwa Peternakan 2.
- Puguh, W., Arhabsi, W. O. R. S., Hastian., 2020. Penambahan Jenis Tepung Terhadap Kualitas Dodol Pisang Mas. Sultra Journal of Agricultural Research, 1(13), 14–25.
- Purba, A.P., Dwiloka, B. Rizqiati, H., 2018. Pengaruh lama fermentasi terhadap bakteri asam laktat (BAL), viskositas, aktivitas antioksidan, dan organoleptik water kefir anggur merah (*Vitis vinifera* L.). J. Teknologi Pangan, 2(1): 49-51.
- Puspitarini, O.R., I. Kentjonowaty., Rasbawati., 2024. Kadar laktosa, total solid, dan solid non fat kefir susu sapi yang diolah dari jenis susu berbeda. J. Peternakan, 21 (1): 33-37.
- Putri, Y. D., Setiani, N. A., Warya, S., 2020. The Effect of Temperature, Incubation and Storage Time on Lactic Acid Content, pH, and Viscosity of Goat Milk Kefir. Current Research on Biosciences and Biotechnology, 2(1), 101–104. <https://doi.org/10.5614/crbb.2020.2.1/hpmq5042>
- Rasbawati, R., Irmayani, I.D. Novieta., Nurmianti, N., 2019. Karakteristik Organoleptik dan Nilai pH Yoghurt dengan Penambahan Sari Buah Mengkudu (*Morinda citrifolia* L). Jurnal Ilmu Produksi Dan Teknologi Hasil Peternakan, 7(1), 41–46 <https://doi.org/10.29244/jipthp.7.1.41-46>
- Rizqiati, H., Ramadhanti, D. L., Prayoga, M. I. Y., 2021. Pengaruh Variasi Konsentrasi Sukrosa Terhadap Total Bakteri Asam Laktat, pH, Kadar Alkohol dan Hedonik Water Kefir Belimbing Manis (*Averrhoa carambola*). Jurnal Ilmiah Sains, 21(1), 54. <https://doi.org/10.35799/jis.21.1.2021.3110>
- Rohmah, F., Estiasih, T., 2018. Perubahan Karakteristik Kefir Selama Penyimpanan : Kajian Pustaka. Jurnal Pangan Dan Agroindustri, 6(3), 30–36. <https://doi.org/10.21776/ub.jpa.2018.006.03.4>
- Rosa, D. D., Dias, M. M. S., Grzeškowiak, Ł. M., Reis, S. A., Conceição, L. L., Peluzio, M. D. C. G., 2017, June 1. Milk kefir: Nutritional, microbiological and health benefits. Nutrition Research Reviews. Cambridge University Press. <https://doi.org/10.1017/S0954422416000275>

- Safitri, M. F., A. Swarastuti., 2013. Kualitas kefir berdasarkan konsentrasi kefir grains. Jurnal Aplikasi Teknologi Pangan. 2 (2) : 87 – 92.
- Savaiano, D.A., Hutkins, R.W., 2021. Yogurt, cultured fermented milk, and health: A systematic review. Nutr Rev 79, 599–614.
- Sinurat, R. L., Ekowati, C. N., Sumardi, S., Farisi, S., 2018. Karakteristik Kefir Susu Sapi dengan Inokulum Ragi Tape. Jurnal Ilmiah Peternakan Terpadu, 6(2), 111–116. <https://doi.org/10.23960/jipt.v6i2.p111-116>
- Ton, A.M.M., Campagnaro, B.P., Alves, G.A., Aires, R., Côco, L.Z., Arpini, C.M., Guerra E Oliveira, T., Campos-Toimil, M., Meyrelles, S.S., Pereira, T.M.C., Vasquez, E.C., 2020. Oxidative Stress and Dementia in Alzheimer's Patients: Effects of Synbiotic Supplementation. Oxid Med Cell Longev 2020, 1–14.
- Tuhumury, H. C. D., Moniharapon, E., Souripet, A., 2018. Karakteristik Sensoris Puree Pisang Tongka Langit Pendek (*Musa troglodytarum*). Jurnal Teknologi Pertanian, 9(2), 1–10.
- Utomo, B., Marsiti, C. I. R., Damiani, D., 2019. Uji Kualitas Tepung Pisang Mas (*Musa Acuminata*). Jurnal BOSAPARIS: Pendidikan Kesejahteraan Keluarga, 9(3), 189. <https://doi.org/10.23887/jjpkk.v9i3.22146>
- Viogenta, P., Kartinah, N., Khairunnisa, A., Rahman, F., 2021. Quality of Peanut (*Arachis hypogaeae* L.) Kefir with Variation in Ragi Starter Concentration and Long Fermentation. Jurnal Biota, 7(2), 85–93. <https://doi.org/10.19109/biota.v7i2.8147>
- Y. Fathi, N. Ghodrati, M.-J. Zibaeenezhad., S. Faghieh., 2017. Kefir drink causes a significant yet similar improvement in serum lipid profile, compared with low-fat milk, in a dairy rich diet in overweight or obese premenopausal women: a randomized controlled trial, Journal of Clinical Lipidology, vol. 11, no. 1, pp. 136–146, 2017.
- Yilmaz, B., Sharma, H., Melekoglu, E., Ozogul, F., 2022. Recent developments in dairy kefir-derived lactic acid bacteria and their health benefits. Food Bioscience, 46. <https://doi.org/10.1016/j.fbio.2022.101592>
- Yousefvand, A., Huang, X., Zarei, M., Saris, P. E. J., 2022. Lacticaseibacillus rhamnosus GG Survival and Quality Parameters in Kefir Produced from Kefir Grains and Natural Kefir Starter Culture. Foods, 11(4). <https://doi.org/10.3390/foods11040523>