Indigenous endomycorrhiza of iron and manganese contaminated soils in Indonesia

by Cek Turnitin


Submission date: 01-May-2023 01:48PM (UTC+0800)

Submission ID: 2080652957

File name: 2._2019_publikasi_Indigenous_endomycorrhiza_of_i.pdf (330.34K)

Word count: 4142

Character count: 23132





International Journal of Current Research in Biosciences and Plant Biology

Volume 6 • Number 8 (August-2019) • ISSN: 2349-8080 (Online)



Journal homepage: www.ijcrbp.com

Original Research Article

doi: https://doi.org/10.20546/ijcrbp.2019.608.001

Indigenous endomycorrhiza of iron and manganese contaminated soils in Indonesia

Muhammad Akhsan Akib¹*, Andi Nuddin¹, Retno Prayudyaningsih² and Syatrawati³

¹Muhar 18 adiyah University of Parepare, South Sulawesi, 91131, Indonesia ²Environment and Forestry Research and Development Institute of Makassar, Indonesia ³Polytechnic of Pangkep State Agricultural, Indonesia

*Corresponding author; e-mail: akhsanbagus@yahoo.co.id

Article Info

Date of Acceptance: 07 July 2019

Date of Publication: 06 August 2019

Keywords

Acaulospora
Fungus
Gigaspora
Heavy metal
Phytorhizoremediation

ABSTRACT

Concentrations of iron and manganese that exceed critical line in soil and plant provide opportunities for microorganisms to become resistant. The aim of this study was to identify endomycorrhizal fungi from areas contaminated with iron and manganese for use as biostarter for phytorizoremediation programs. This research was conducted in two phases at April 2019, namely taking rhizosphere from *Chromolaena odorata*, *Melastoma affine dan Spathoglottis plicata* in Sorowako, Indonesia; and the second phase is to isolate and identify endomycorrhizal fungal spores in the Microbiology Laboratory, Center for Environmental and Forestry Research and Development in Makassar, Indonesia. The results showed that the genus *Acaulospora* was more dominant in areas that were contaminated with iron and manganese, and were able to adapt and survive compared to other genera.

Introduction

Endomycorrhizal a form of mutualism symbiotic association between root endodermis of higher plants with fungal hyphae, which have reached 80% of plant species. Some researchers have shown the positive role of endomycorrhizal on plant growth in stressful environments of salinity (Hashem et al., 7018; Hashem et al., 2019), nutrients (Chanda et al., 2014; Bowles et al., 2018) drought (Abdelmoneim et al., 2013; He et al., 2019), pathogens (Whipps, 2011; Adamec and Andrejiova, 2018) and heavy metals (Miransari, 2011; Atakan et al., 2018).

Research on the use of endomycorrhizal to reduce heavy metal contamination is a phenomenon that has received the attention of phytorhizoremediation researchers. Arisusanti and Purwani (2013) infect Glomus fasciculatum on the roots of Dahlia sp. to reduce the metal, lead (Pb). Rhizophagus clarus and R. irregularis can provide protective effects on vines on land contaminated with copper (Cu) (Ambrosini et al., 2015). Glomus versiforme and Rhizophagus intraradices can reduce the 15 sorption of cadmium (Cd) on Lonicera japonica (Jiang et al., 2016). Alam et al. (2019) and Singh et al. (2015) using endomy 11 hizal species obtained from International (Vesicular)

Arbuscular Mycorrhizal Fungi (INVAM), West Virginia University (WV), USA, inoculated on *Lens culinaris* to reduce arsenic (As), but the endomycorrhizal used are not indigenous endomycorrhizal from areas that contaminated with heavy metals.

The chemical elements iron (Fe) and manganese (13 n), respectively, have specific gravity of 7.86 g.cm⁻³ and 7.21 g.cm⁻³, so that classified as heavy metals (Engge and Hendrajaya, 2016; Dangeand Dhoble, 2017). Fe and Mn including as microessential heavy metals that can cause metabolic disorders for soil microorganisms if are in high concentrations (Singh and Prasad, 2015; Chu, 2017). Some research results to show that heavy metal polition give out detrimental effect on the life cycle of soil microorganisms (Xie et al., 2016; Jin et al., 2018) and causes changes in the soil microoganism community structure (Gurave et al., 2015; Wood et al., 2016, Park et al., 2016), but ultimately causes soil microor nisms to become tolerant and resistant (Gurave et al, 2015; Harman and Uphoff, 2019). However, the level of tolerance and resistance between groups of fungus is largely determined by the type of fungus and environment (Ramage et al., 2012; Domka et al., 2019).

Researchers have found several strains of endomycorrhizal fungus that are tolerant of heavy metal stress, namely, **Ggaspora albida** and **Glomus clarum** tolerant of zinc (Zn), cadmium (Cd), lead (Pb) and copper (Cu) (Soares and Siqueira, 2008), **Glomus intraradiaces** and **Glomus mosseae** tolerant of Pb and Cd (Adewole et al., 2010; Aram and Golchin, 2013), **Acaulospora morrowiae** tolerant of Cu (Ambrosini et al., 2015), **Acaulospora mellea** tolerant to nickel (Ni) and chromium (Cr) (Spruyt et al., 2014) and **Scutellospora pellucida** tolerant of Zn (Filho et al., 2002; Bano and Ashfaq, 2013).

Endomycorrhizal tolerant and resistant to heavy metals with high concentrations can be a biotechnology agent for the success of post-mining land phytorhizoremediation programs in Sorowako, Indonesia. So that exploring the indigenous Endomycorrhizal fungi spores from areas contaminated with Fe and Mn is the objective of this study, which also provides an opportunity for others researcher to inoculate on endemic plant of location.

Materials and methods

The study conducted in two phases at April 2019. The first phase, rhizosphere collection of *Chromolaena odorata*, *Melastoma affine* and *Spathoglottis plicata* at coordinated of 2°31'57.6 "S and 121°22'50.7" E, Sorowako, Indonesia. The rhizosphere collection method uses a diagonal system (Subramani, 2009; Astiko et al, 2016).

The second phase, endomycorrhizal fungus spores isolated from rhizosphere using the wet sieving technique (Chanda et al., 2014; Shamini and Amutha, 2014) and the sucrose gradient centrifugation method (Walker et al., 1982; Charoenpak et al., 2010) at Microbiology Laboratory, Research and Development Center for Environment and Forestry, Makassar, Indonesia. Spore 3 orphology identified using a manual book from the International Culture Collection of Vesicular Arbuscular Mycorrhizal Fungus (INVAM, 2019).

Concentration Fe and Mn in soil was measured while in laboratory of chemistry, Polytechnic of Ujung Pandang, Makassar, using X-Ray Florence Spectrophotometer/Bruker/S2 Ranger, concentration Fe and Mn in soil can be seen in Fig. 1.

Results and discussion

Concentrations of Fe and Mn in nickel post-mine area of Sorowako have exceeded critical limit for the environment and plants (Fig. 1), some researchers suggest that critical limit for Fe in the soil were 100,000 ppm (Mengel 2nd Kirdy, 1987) and in plants were 1,000 ppm (Ministry of State for Population and Environment of the Republic of Indonesia and Dalhousie University Canada, 1992), while for Mn, the critical limit in the soil were 1,500 ppm (Liphadzi and Kirkham, 2005) and in plants was 400 ppm (Jones, 1979).

Fe concentration and high Mn is likely caused by mining activities, especially when returning overburden, so that it can have a toxic effect on the activity of soil microorganisms to develop and reproduce, but according to Ivshina et al (2013) that strategy to adaptation and tolerance also owned by each organism in unfavorable environment.

Endomycorrhizal also have defense strategy to heavy metal stress, the strategy is likely to involve one of the mechanisms, 19 cluding: (i) new expression of fungus gene (Cicatelli et al., 2010; Ferrol et al., 2016), (ii) metal of quarantined and deposited at extracellular (Koohi, 2014; Chen et al., 2018), (iii) produce of metal binding protein (Singh, 2012; Hrynkiewicz et al., 2012), (iv) reduce of metal absention (Karimi et al., 2011; Emamverdian et al., 2015), (v) increase efflux (Hildebrandt et al., 2007; Tiwari and Lata, 2018), (vi) formation of complexes outside the cell (Violante et al., 2010; Singh et al. 22016), (vii) release of organic acids (Adeleke et al., 2017; Mishra et al., 2017), and (viii) ligand synthesis such as polyphosphates and metallothionein (Ferrol et al., 2016; Zhan et al., 2019).




Fig. 1: Concentrations of Fe and Mn in nickel postmine area of Sorowako, Indonesia (CLS, Critical limit in soil; CLP, Critical limit in plant). Data is displayed in logarithmic form.

Indigenous endomycorrhiza spores identified in various plant rhizosphere, obtained spores from two genera that are able to adapt in areas that have high concentrations of Fe and Mn, namely *Acaulospora sp.* and *Gigaspora sp.*; *Acaulospora sp.* was the genus mycorrhiza which belongs to the family Acaulosporaceae. This genus has several characteristics including having 2-3 spore walls, spores formed on the side of the neck of the sporiferoussaccule, globose to elliptical, hyaline, yellow, or yellowish red, spore diameter between 74-289 μm (INVAM, 2019), however, the diameter of the spores found by Akib et al. (2018) on nickel-contaminated land ranged from 60 - 80 μm.

Gigaspora sp. was the mycorrhizal genus that belongs to the Gigasporaceae family. This genus has characteristics, among others, spores are produced singly in the soil, not having a layer of inner spore walls, have bulbous suspensors, globose or sub-globose shaped, creamy to yellow in color, 206-358 μ m in diameter (INVAM, 2019). The recearch results of Akib et al. (2018) in the nickel post-mine area found spores of Gigaspora sp. with diameter of 203 - 235 μ m.




Fig. 2: Spores morphology of *Acaulospora sp.* (a) and *Gigaspora sp.* (b) which isolated from area contaminated with Fe and Mn (Note: sw, cell wall; sc, saccule; sh, substanding hyphae; hf, hyphae).

The presence of indigenous endomycorrhizal spores in environment contaminated with Fe and Mn, allegedly do one of strategic defense, as has explained by researchers. However, according to Herrera et al. (2018), the ability of adaptation and resistance of indigenous endomycorrhizal to be tolerant, it is possible follow the intracellular metal binding mechanism through ligand synthesis (metallothionein, polyphosphate), and / or accumulate metals to in vacuoles.

Calculation of spore's indigenous endomycorrhizal number per 1000 mg of rhizosphere contaminated with Fe and Mn were found in different amounts, and dominated by the genus *Acaulospora sp.* (Table 1). This is probably due to *Acaulospora sp.* having more than one cell wall, so it has a stronger defense against heavy metal stress than *Gigaspora sp.* According to Upadhyaya et al. (2010) and Koohi (2014) that endomycorrhizae can store heavy metals in hyphal cell walls, specifically stored in crystalloids in the 12 ycelium, this explanation reinforced by Chaves et al. (2002) and Chen et al. (2018) who show that in roots of plants that are contaminated with heavy metals, it is seen

that the endomycorrhizal extra radical mycelium is able to absorb and accumzoate heavy metals outside the mushyel wall of hyphal wall zone, in cell wall, and in cytoplasm of fungus hyphae, while the research results of Tuheteru et al. (2017) show that utilization *Acaulospora tuberculata* can increase Mn, Fe, Cr and Ni uptake of *Nauclea orientalis* L.

Table 1. Number of indigenous endomycorrhizal spores per 1000 mg of rhizosphere sample.

Family	Rhizosphere of pioneer plant	Number of spores	
		GS	AC
Asteraceae	Chromolaena odorata	О	24
Melastomataceae	Melastoma affine	1	1
Nephrolepidaceae	Nephrolepis exaltata	О	13

Note: GS, Gigaspora sp.; AC, Acaulospora sp.

Several studies have reported that indigenous endomycorrhiza effectively and efficiently infect endodermis of plant roots in areas that are contaminated with heavy metals (Vaishaly et al., 2015; Shakeel and Yaseen, 2015; Kumar and Saxena, 2019). According to Zadehbagheri et al. (2014) that indigenous endomycorrhizal isolates that develop and reproduce naturally in areas contaminated with heavy metals are more tolerant than isolates originating from non-polluted areas. filtering tolerant indigenous Thus, endomycorrhizal isolates needs to be done to ensure the effectiveness of symbiosis between endomycorrhiza and endodermis of plant roots. Therefore, it is very important to infect endodermis of endemic plant roots with indigenous endomycorrhizal isolates in future studies.

Conclusion

Indigenous endomycorrhizal of genus *Acaulospora sp.* and *Gigaspora sp.* that are able to adapt in areas contaminated with Fe and Mn have been found in Sorowako and can be used as sources of inoculum in phytorhizoremediation programs combined with endemic plants.

Acknowledgement

Authors sincerely thank Ministry of Research, Technology and Higher Education of the Republic of Indonesia for providing support through a grant of basic research competition in 2019 and special thanks to Makole Nuha, Mr. H. Andi Baso, AM Opu To Lamattulia, my brother Mr. Syamsul Amri Akib, Mr. Komaruddin and friends, for all the help during our activities in Sorowako.

Conflict of interest statement

Authors declare that they have no conflict of interest.

References

Abdelmoneim, T. S., Moussa, T. A. A., Almaghrabi, O. A., Alzahrani, H. S., Abdelbagi, I., 2013. Increasing plant tolerance to drought Stress by inoculation with arbuscular mycorrhizal fungi. Life Sci. J. 10, 3273-3280.

Adamec, S., Andrejiova, A., 2018. Mycorrhiza and stress tolerance of vegetables. Acta Hortic. Regiotec. 2, 30-35.

Adeleke, R., Nwangburuka, C., Oboiriende, B., 2017. Origins, roles and fate of organic acids in soils. South Afr. J. Bot. 108, 393-406.

Adewole, M. B., Awotoye, O. O., Ohiembor, M. O., Salami, A. O., 2010. Influence of mycorrhizal fungi on phytoremediating potential and yield of sunflower in Cd and Pb polluted soils. J. Agric. Sci. 55, 17-28.

Akib, M. A., Mustari, K., Kuswinanti, T., Syaiful, S, A., 2018. Identification and abundance of indigenous endomycorrhiza isolated from nickel post-mining plantation in Sorowako. Int. J. Curr. Res. Biosci. Plant Biol. 5(4), 8-16.

Alam, M. Z., Hoque, M. A., Ahammed, G. J., Carpenter-Boggs, L. 2019. Arbuscular mycorrhizal fungi reduce arsenic uptake and improve plant growth in *Lens culinaris*. Plos One 14(5), e0211441.

Ambrosini, V. G., Voges, J. G., Canton, L., Couto, R. R., Ferreira, P. A. A., Comin, J. C., Melo, G. W. B., Brunetto, G., Soares, C. R. F. S., 2015. Effect of arbuscular mycorrhizal fungi on young vines in copper-contaminated soil. Braz.

- J. Microbiol. 46, 1045-1052.
- Aram, H., and Golchin, A. 2013. The effects of arbuscular mycorrhizal fungi on nitrogen concentration of berseem clover in contaminated soil with cadmium. J. Chem. Health Risks 3, 53-58.
- Arisusanti, R. J., Purwani, K. I., 2013. Pengaruh Mikoriza *Glomus fasciculatum* terhadap Akumulasi Logam Timbal (Pb) pada Tanaman *Dahlia pinnata*. J. Sains SeniPomits 2, 69-73.
- Astiko, W., Fauzi, M. T., Sukartono, 2016. Mycorrhizal population on various cropping systems on sandy soil in Dryland Area of North Lombok, Indonesia. Nusantara Biosci. 8, 66-70.
- Atakan, A., Ozkaya, H. O., Erdogan, O. 2018. Effects of arbuscular mycorrhizal fungi (AMF) on heavy metal and salt stress. Turk. J. Agric. Food Sci. Technol. 6, 1569-1574.
- Bano, S. A., Ashfaq, D., 2013. Role of mycorrhiza to reduce heavy metal stress. Natural Sci. 5, 16-20.
- Bowles, T. M., Jackson, L. E., Cavagnaro, T. R., 2018. Mycorrhizal fungi enhance plant nutrient acquisition and modulate nitrogen loss with variable water regimes. Glob. Chang. Biol. 24, e171-e182.
- Chanda, D., Sharma, G. D., Jha, D. K., 2014. Isolation and identification of some arbuscular mycorrhiza (AM) fungi for phytoremediation in soil contaminated with paper mill effluent. Int. J. Curr. Microbiol. App. Sci. 3, 527-539.
- Charoenpakdee, S., Phosri, C., Dell, B., Lumyong, S., 2010. The mycorrhizal status of indigenous arbuscular mycorrhizal fungi of physic nut (*Jatropha curcas*) in Thailand. Mycosphere 1, 167–181.
- Chavez, C. G., D'Haen, J., Vangronsveld, J., Dodd, J. C. 2002. Copper sorption and accumulation by the extraradical mycelium of different *Glomus spp*. (arbuscular mycorrhizal fungi) isolated from the same polluted soil. Plant Soil 240, 287–297.
- Chen, B., Nayuki, K., Kuga, Y., Zhang, X., Wu, S., Ohtomo, R., 2018. Uptake and intraradical immobilization of cadmium by arbuscular mycorrhizal fungi as revealed by a stable isotope tracer and synchrotron radiation μXray fluorescence analysis. Microbes Environ. 33, 257-263.
- Chen, M., Arato, M., Borghi, L., Nouri, E., Reinhardt, D., 2018. Beneficial services of

- arbuscularmycorrhizal fungi from ecology to application. Front. Plant Sci. 9, 1270.
- Chu, D., 2017. Effects of heavy metals on soil microbial community. IOP Conf. Series: Earth and Environmental Science 113: 012009.
- Cicatelli, A., Lingua, G., Todeschini, V., Biondi, S., Torrigiani, P., Castiglione, S., 2010. Arbuscular mycorrhizal fungi restore normal growth in a white poplar clone grown on heavy metal-contaminated soil, and this is associated with upregulation of foliar metallothionein and polyamine biosynthetic gene expression. Ann. Bot. 106, 791-802.
- Dange, S. S., Dhoble, R. M., 2017. Iron removal by adsorption. Int. J. Sci. Technol. Eng. 3, 603-606.
- Domka, A. M., Rozpadek, P., Turnau, K., 2019. Are fungal endophytes merely mycorrhizal copycats? the role of fungal endophytes in the adaptation of plants to metal toxicity. Front. Microbiol. 10, 371.
- Emamverdian, A., Ding, Y., Mokhberdoran, F., Xie, Y., 2015. Heavy metal stress and some mechanisms of plant defense response. Scient. World J. Article ID 756120. 18 pages.
- Engge, Y., Hendrajaya, L., 2016. Prinsip Dan Proses Fisika Dalam Penambangan Mangan (Mn) Di Kabupaten Timor Tengah Selatan, Provinsi Nusa Tenggara Timur. Prosiding Seminar Nasional Pendidikandan Saintek, pp.286-296.
- Ferrol, N., Tamayo, E., Vargas, P., 2016. The heavy metal paradox in arbuscular mycorrhizas: From mechanisms to biotechnological applications. J. Exp. Bot. 67, 6253-6265.
- Filho, O. K., Siqueira, J. O., Moreira, F. M. S., 2002. Vesicular arbuscular mycorrhizae fungi in soils polluted with heavy metals. Rev. Bras. Cien. Solo 26, 125-134.
- Gurave, N. A., Korde, V. V., Dhas, S. S., Disale, M., 2015. Isolation and identification of heavy metal resistant bacteria from petroleum soil of Loni, Ahmednagar. Eur. J. Exp. Biol. 5, 6-11.
- Harman, G. E., Uphoff, N., 2019. Symbiotic rootendophytic soil microbes improve crop productivity and provide environmental benefits. Hindawi Scientifica, Article ID 9106395.
- Hashem, A., Abd_Allah, E. F., Alqarawi, A. A., Wirth, S., Egamberdieva, D., 2019. Comparing symbiotic performance and physiological responses of two soybean cultivars to

- arbuscular mycorrhizal fungi under salt stress. Saudi J. Biol. Sci. 26, 38-48.
- Hashem, A., Alqarawi, A. A., Radhakrishnan, R., Al-Arjani, A. F., Aldehaish, H. A., Egamberdieva, D., Abd_Allah, E. F., 2018. Arbuscular mycorrhizal fungi regulate the oxidative system, hormones and ionic equilibrium to trigger salt stress tolerance in *Cucumis sativus* L. Saudi J. Biol. Sci. 25, 1102–1114.
- He, J. D., Wu, Q. S., Zou, Y. N., 2019. Effects of mycorrhiza and drought stress on the diversity of fungal community in soils and roots of trifoliate orange. Biotechnology 18, 32-41.
- Herrera, H., Valadares, R., Oliveira, G., Fuentes,
 A., Almonacid, L., Nascimento, S. V., Bashan,
 Y., Arriagada, C., 2018. Adaptation and
 tolerance mechanisms developed by
 mycorrhizal *Bipinnula fimbriata* plantlets
 (Orchidaceae) in a heavy metal-polluted
 ecosystem. Mycorrhiza 28, 651.
- Hildebrandt, U., Regvar, M., Bothe, H., 2007. Arbuscular mycorrhiza and heavy metal tolerance. Phytochemistry 68, 139-146.
- Hrynkiewicz, K., Dabrowska, G., Baum, C., Niedojadlo, K., Leinweber, P. 2012. interactive and single effects of ectomycorrhiza formation and bacillus cereus on metallothionein mt1 expression and phytoextraction of cd and zn by willows. Water Air Soil Pollut. 223, 957-968.
- INVAM, 2019. International Culture Collection of Vesicular ArbuscularMycorrhizal Fungi. http://invam.caf.wvu.edu
- Ivshina, I. B., Kuyukina, M. S., Kostina, L. V. 2013. Adaptive mechanisms of nonspecific resistance to heavy metal ions in alkanotrophic actinobacteria. Russ. J. Ecol. 44, 123-130.
- Jiang, Q. Y., Zhuo, F., Long, S. H., Zhao, H. D., Yang, D. J., Ye, Z. H., Li, S. S., Jing, Y. X., 2016. Can arbuscular mycorrhizal fungi reduce Cd uptake and alleviate Cd toxicity of *Lonicera japonica* grown in Cd-added soils? Scient. Rep. 6, 21805.
- Jin, Y., Luan, Y., Ning, Y., Wang, L., 2018. Effects and mechanisms of microbial remediation of heavy metals in soil: A critical review. Appl. Sci. 8, 1336.
- Jones, U.S., 1979. Fertilizers and Soil Fertility. Reston Publ. Co., Virginia. 264p.
- Karimi, A., Khodaverdiloo, H., Sepehri, M., Sadaghiani, M. S., 2011. Arbuscular mycorrhizal fungi and heavy metal

- contaminated soils. Afr. J. Microbiol. Res. 5, 1571-1576.
- Koohi, S. S., 2014. Role of arbuscular mycorrhizal (AM) fungi in phytoremediation of soils contaminated: a review. Int. J. Adv. Biol. Biomed. Res. 2, 1854-1864.
- Kumar, S., Saxena, S., 2019. Arbuscular mycorrhizal fungi (AMF) from heavy metal-contaminated soils: molecular approach and application in phytoremediation. In: Biofertilizers for Sustainable Agriculture and Environment (Eds.: Giri, B., Prasad, R., Wu, Q.S., Varma, A.). Soil Biol. 55, 489-500.
- Liphadzi, M. S., Kirkham, K. B., 2005. Phytoremediation of soil contaminated with heavy metals: a technology for rehabilitation of the environment. South Afr. J. Bot. 71(1), 24-37.
- Mengel, K., Kirby, E., 1987. Principles of Plant Nutrition. International Potash Inst. Bern Switzerland.
- Ministry of State for Population and Environment Republic of Indonesia and Dalhousie University Canada. 1992. Environmental Management in Indonesia. Report on Soil Quality Standards for Indonesia (Interim Report).
- Miransari, M., 2011. Hyperaccumulators, arbuscular mycorrhizal fungi and stress of heavy metals. Biotechnol. Adv. 29, 645-653.
- Mishra, J., Singh, R., Arora, R. K., 2017. Alleviation of heavy metal stress in plants and remediation of soil by rhizosphere microorganisms. Front. Microbiol. 8, 1706.
- Park, H., Lee, E. H., Ka, K. H., Eom, A. H. 2016. Community structures of arbuscular mycorrhizal fungi in soils and plant roots inhabiting abandoned mines of Korea. Mycobiology 44, 277-282.
- Ramage, G., Rajendran, R., Sherry, L., Williams, C., 2012. Fungal biofilm resistance. Int. J. Microbiol., Article ID 528521.
- Shakeel, M., Yaseen, T., 2015. An insight into phytoremediation of heavy metals from soil assisted by ancient fungi from Glomeromycotaarbuscular mycorrhizal fungi. Sci. Technol. Develop. 34, 215-220.
- Shamini, S., Amutha, K. D., 2014. Techniques for extraction of arbuscular mycorrhizal fungi spores. Int. Front. Sci. Technol. 2, 1-6.
- Singh, A., Prasad, S. M., 2015. Remediation of heavy metal contaminated ecosystem: An

- overview on technology advancement. Int. J. Environ. Sci. Technol. 12, 353-366.
- Singh, P. K., 2012. Role of glomalin related soil protein produced by arbuscular mycorrhizal fungi: A review. Agric. Sci. Res. J. 2, 119-125.
- Singh, R., Singh, S., Parihar, P., Singh, V. P., Prasad, S. M., 2015. Arsenic contamination, consequences and remediation techniques: a review. Ecotoxicol. Environ. Saf. 112, 247-270.
- Singh, S., Parihar, P., Singh, R., Singh, V. P., Prasad, S. M., 2016. Heavy metal tolerance in plants: Role of transcriptomics, proteomics, metabolomics, and ionomics. Front. Plant Sci. 6, 1143.
- Soares, C. R. F. S., Siqueira, J. O., 2008. Mycorrhiza and phosphate protection of tropical grass species against heavy metal toxicity in multi-contaminated soil. Biol. Fertil. Soils, 44, 833-841.
- Spruyt, A., Buck, M. T., Mia, A., Straker, C. J., 2014. Arbuscular mycorrhiza (AM) status of rehabilitation plants of mine wastes in South Africa and determination of AM fungal diversity by analysis of the small subunit rRNA gene sequences. South Afr. J. Bot. 94, 231-237.
- Subramani, J., 2009. Further results on diagonal systematic sampling scheme for finite populations. J. Indian Soc. Agric. Stat. 63, 277-282.
- Tiwari, S., Lata, C., 2018. Heavy metal stress, signaling, and tolerance due to plant-associated microbes. Front Plant Sci. 9, 452.
- Tuheteru, F. D., Arif, A., Widiastuti, E., Rahmawati, N., 2017. Heavy metal uptake by indigenous arbuscular mycorrhizas of *Nauclea orientalis* L. and the potential for phytoremediation of serpentine soil. J. Ilmu Kehutanan 11, 76-84.
- Upadhyaya, H., Panda, S. K., Bhattacharjee, M. K., and Dutta, S. 2010. Role of arbuscular

- mycorrhiza in heavy metal tolerance in plants: Prospects for phytoremidiation. J. Phytol. 2, 16-27.
- Vaishaly, A. G., Mathew, B. B., Krishnamurthy, N. B., Krishnamurthy, T. P., 2015. Bioaccumulation of heavy metals by fungi. Int. J. Environ. Chem. Chromatogr. 1, 15-21.
- Violante, A., Cozzolino, V., Perelomov, L., Caporale, A. G., Pigna, M., 2010. Mobility and bioavailability of heavy metals and metalloids in soil environments. J. Soil. Sci. Plant Nutr. 10: 268 292.
- Walker, C., Mize, C. W., McnabbJr, H. S., 1982.Population of endogonaceus at two locationin central Iowa. Can. J. Bot. 60, 2518-2529.
- Whipps, J. M., 2011. Prospects and limitations for mycorrhizas in biocontrol of root pathogens. Can. J. Bot. 82, 1198-1227.
- Wood, J. L., Zhang, C., Mathews, E. R., Tang, C., Franks, A. E., 2016. Microbial community dynamics in the rhizosphere of a cadmium hyper-accumulator. Scient. Rep. 6, 36067.
- Xie, Y., Fan, J., Zhu, W., Amombo, E., Lou, Y., Chen, L., Fu, J., 2016. Effect of heavy metals pollution on soil microbial diversity and Bermudagrass genetic variation. Front. Plant Sci. 7, 755.
- Zadehbagheri, M., Azarpanah, A., Javanmardi, S., 2014. Perspective of arbuscular mycorrhizal fungi phytoremediation on contamination and remediation heavy metals soil in sustainable agriculture. Amer.-Eur. J. Agric. Environ. Sci. 14, 379-386.
- Zhan, F., Li, B., Jiang, M., Li T., He Y., Li, Y., Wang, Y., 2019. Effects of arbuscular mycorrhizal fungi on the growth and heavy metal accumulation of Bermudagrass [Cynodon dactylon (L.) Pers.] grown in a leadzinc mine wasteland. Int. J. Phytoremed. 21,

How to cite this article:

Akib, M. A., Nuddin, A., Prayudyaningsih, R., Syatrawati, 2019. Indigenous endomycorrhiza of iron and manganese contaminated soils in Indonesia. Int. J. Curr. Res. Biosci. Plant Biol. 6(8), 1-7. **doi:** https://doi.org/10.20546/ijcrbp.2019.608.001

Indigenous endomycorrhiza of iron and manganese contaminated soils in Indonesia

ORIGINALITY REPORT

1 %
SIMILARITY INDEX

5%
INTERNET SOURCES

9%
PUBLICATIONS

U% STUDENT PAPERS

PRIMARY SOURCES

Asma Fayyad Bzour, Hani Nicola Khoury, Sawsan Attalah Oran. "Uptake of Some Toxic Elements by Wild Plants in Siwaqa Area/Central Jordan", Applied Microscopy, 2017

%

Publication

Eka Djatnika Nugraha, June Mellawati, Wahyudi, Chutima Kranrod et al. "Heavy Metal Assessments of Soil Samples from a High Natural Background Radiation Area, Indonesia", Toxics, 2022

<1%

Publication

M A Akib, K Mustari, T Kuswinanti, S A Syaiful. "Abundance of arbuscular mychorrizal fungi in rehabilitation area of nickel post-mining land of Sorowako, South Sulawesi", IOP Conference Series: Earth and Environmental Science, 2018

<1%

jglitrop.ui.ac.id

<1%

5	mail.scialert.net Internet Source	<1%
6	pubmed.ncbi.nlm.nih.gov Internet Source	<1%
7	real-phd.mtak.hu Internet Source	<1%
8	ur.booksc.me Internet Source	<1%
9	www.i-scholar.in Internet Source	<1%
10	Kendall R. Corbin, Bridget Bolt, Carlos M. Rodríguez López. "Breeding for Beneficial Microbial Communities Using Epigenomics", Frontiers in Microbiology, 2020 Publication	<1%
11	Mominur Rahman, Ashraf Hossain, Mohammad Eyakub Ali, Faridul Alam, Khairul Alam. " Arbuscular mycorrhizal colonization increases yield of mungbean (L.) at moderate level of salinity ", Soil Science and Plant Nutrition, 2019 Publication	<1%
12	Songlin Wu, Xin Zhang, Longbin Huang, Baodong Chen. "Arbuscular mycorrhiza and plant chromium tolerance", Soil Ecology	<1%

Letters, 2019

13	www.diva-portal.org Internet Source	<1%
14	www.fao.org Internet Source	<1%
15	Jia-Dong He, Ge-Ge Chi, Ying-Ning Zou, Bo Shu, Qiang-Sheng Wu, A.K. Srivastava, Kamil Kuča. "Contribution of glomalin-related soil proteins to soil organic carbon in trifoliate orange", Applied Soil Ecology, 2020 Publication	<1%
16	biblio.ugent.be Internet Source	<1%
17	helda.helsinki.fi Internet Source	<1%
18	jurnal.balithutmakassar.org Internet Source	<1%
19	Joske Ruytinx, Arjun Kafle, Muhammad Usman, Laura Coninx, Sabine D. Zimmermann, Kevin Garcia. "Micronutrient transport in mycorrhizal symbiosis; zinc steals the show", Fungal Biology Reviews, 2020 Publication	<1%
20	Philippe Giasson. "Arbuscular Mycorrhizae and Alleviation of Soil Stresses on Plant	<1%

Growth", Mycorrhizae Sustainable Agriculture and Forestry, 2008

Publication

Renata Kurtyka, Zbigniew Burdach, Agnieszka Siemieniuk, Waldemar Karcz. "Single and combined effects of Cd and Pb on the growth, medium pH, membrane potential and metal contents in maize (Zea mays L.) coleoptile segments", Ecotoxicology and Environmental Safety, 2018

<1%

- Publication
- Shalini Tiwari, Charu Lata. "Heavy Metal Stress, Signaling, and Tolerance Due to Plant-Associated Microbes: An Overview", Frontiers in Plant Science, 2018

<1%

- Publication
- Ilika Ghosh, Manosij Ghosh, Anita Mukherjee.
 "Chapter 7 Remediation of Mine Tailings and
 Fly Ash Dumpsites: Role of Poaceae Family
 Members and Aromatic Grasses", Springer
 Science and Business Media LLC, 2017

<1%

Publication

Exclude quotes

On

Exclude matches

Off