

20. Bio-Conferences 2024.pdf



Document Details

Submission ID

trn:oid:::3618:96695964

Submission Date

May 19, 2025, 8:07 PM GMT+7

Download Date

May 19, 2025, 8:09 PM GMT+7

File Name

20. Bio-Conferences 2024.pdf

File Size

592.2 KB

13 Pages**5,469 Words****29,935 Characters**

20% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.





Filtered from the Report

- Bibliography
- Quoted Text
- Cited Text
- Submitted works
- Internet sources




Exclusions

- 2 Excluded Sources

Match Groups

-  **67 Not Cited or Quoted 20%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 0%  Internet sources
- 20%  Publications
- 0%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 67 Not Cited or Quoted 20%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 0% Internet sources
- 20% Publications
- 0% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Publication	Idin Sahidin, Baru Sadarun, Adryan Fristiohady, Nur Syifa Rahmatika et al. " Che...	3%
2	Publication	R Rahmi, A Syarif, J Ibrahim, F Indah Yani, A Chadijah, A R Nanda, A S Syamsuri, F ...	2%
3	Publication	Khalid Amjad, Hans-Uwe Dahms, Chuan-Hsin Ho, Yu-Chen Wu, Feng-You Lin, Hon...	1%
4	Publication	Muhammad Iqbal, Tom Amey, Gusti Ayu Widayanti, Ina Aprillia, Randi Syafutra, D...	1%
5	Publication	Ita Apriani, Iis Diatin, Tatag Budiardi, Yani Hadiroseyani, Widanarni Widanarni. "P...	<1%
6	Publication	Muhammed Jefri, Woro Hastuti Satyantini, Adriana Monica Sahidu, Daruti Dinda ...	<1%
7	Publication	"Biotechnological Advances in Aquaculture Health Management", Springer Scienc...	<1%
8	Publication	Arturo Susarrey-Arce, José Federico Hernández-Sánchez, Marco Marcello, Yuri Dia...	<1%
9	Publication	Iko Imelda Arisa, Julianda Fatma, Nurfadillah Nurfadillah, Cut Nuzlia, Ulfa Rahmi. ...	<1%
10	Publication	Muhammad Fachri, Kwaku Amoah, Yu Huang, Jia Cai et al. "Probiotics and parapr...	<1%

11	Publication	Ghasem Mohammadi, Gholamreza Rafiee, Mohammed F. El Basuni, Hien Van Do...	<1%
12	Publication	Julatat Keereelang, Kriangsak Mangumphan, Chanagun Chitmanat, Sudaporn To...	<1%
13	Publication	D.P Hartono, E Barades. "Effectiveness of using Commercial Probiotics in Biofloc S...	<1%
14	Publication	Ricky Febrinaldy Simanjuntak, Ira Maya Abdiani, Perdiansyah Perdiansyah, Riska ...	<1%
15	Publication	Dony Prasetyo, Anis Zubaidah, Rachardian Dwi Cahya Putra, Olga Anne, Firdaus A...	<1%
16	Publication	Vini Taru Febriani Prajayati, Otie Dylan Subhakti Hasan, Mugi Mulyono. "The Dige...	<1%
17	Publication	Sumitro Sumitro, Arfan Afandi, Wa Ode Safia. "Evaluation of Flock Volume Levels ...	<1%
18	Publication	Angela Mariana Lusiastuti, Sari Dwi Maryanti, Uni Purwaningsih Purwaningsih. "P...	<1%
19	Publication	B.K. Dey, M.C.J. Verdegem, M.A.J. Nederlof, K. Masagounder, J. Mas-Muñoz, J.W. Sc...	<1%
20	Publication	Jun-Young Bae, Gun Hyun Park, Jeong-Yeol Lee, Okorie Eme Okorie, Sungchul C. B...	<1%
21	Publication	"Novel Approaches Toward Sustainable Tilapia Aquaculture", Springer Science an...	<1%
22	Publication	Daniel M. Aguiar, Thayza F. Ziliani, Xiaofeng Zhang, Andreia L.T. Melo et al. "A nov...	<1%
23	Publication	Kajal Kumari, Sujata Sahoo, Sapana Devi Khumujam, Samikshya Mishra et al. "Eff...	<1%
24	Publication	Maftuch, N D A Wulan, H Suprastyani, E Wijayanto, M Noercholis, A A Prihanto, A ...	<1%

25	Publication	S P Febri, A Fikri, Hanisah, S Nazlia, Zuraidah, A Putriningtias, T M Faisal, Murdhi...	<1%
26	Publication	Santika Anggrahini, Irkham Widiyono, Zein Ahmad Baihaqi, Ahmad Sofyan et al. "...	<1%
27	Publication	Adang Saputra, Fia Sri Mumpuni, Eri Setiadi, Irwan Dwi Setiawan. "KINERJA PERTU...	<1%
28	Publication	Alison K. Gee. "Signatures of Restoration and Management Changes in the Water ...	<1%
29	Publication	Anjan Mukherjee, Goutam Chandra, Koushik Ghosh. "Single or conjoint applicatio...	<1%
30	Publication	Meike Imran, Mulis Mulis, Arafik Lamadi. "PEMBERIAN PROBIOTIK DENGAN DOSI...	<1%
31	Publication	Milad Adel, Reza Safari, Reza Pourgholam, Jalil Zorriehzahra, Maria Ángeles Esteb...	<1%
32	Publication	Muhamad Firdaus Syahmi Sam-on, Shuhaimi Mustafa, Amalia Mohd Hashim, Ah...	<1%
33	Publication	Viola Nur Maghribfa, Putri Desi Wulan Sari, Lailatul Lutfiyah. "Secondary Stress ...	<1%
34	Publication	Wenlong Xu, Xiaoman Hu, Hao Li, Xiangrong Tian, Zhihang Ouyang, Yang Du, Jion...	<1%
35	Publication	Ayja Khayrurraja, Roffi Grandiosa, Fittrie Meyllianawaty Pratiwy, Iskandar Iskand...	<1%
36	Publication	Che-Chun Yi, Chun-Hung Liu, Kuo-Pin Chuang, Yi-Ting Chang, Shao-Yang Hu. "A p...	<1%
37	Publication	Fariq Azhar, Nanang Satria Sumarjan, Siti Hilyana. "Combination of Moringa Leaf ...	<1%
38	Publication	Muhammad A.B. Siddik, Janet Howieson, S.M. Majharul Islam, Ravi Fotedar. "Synb...	<1%

39	Publication	Sai-Wei Chen, Chun-Hung Liu, Shao-Yang Hu. "Dietary administration of probiotic ...	<1%
40	Publication	Sajjad Syed Raffic Ali, Kondusamy Ambasankar, Mohamed Saiyad Musthafa, Ram...	<1%
41	Publication	Shady H. Al-Deriny, Mahmoud A.O. Dawood, Attia A. Abou Zaid, Wael F. El-Tras et ...	<1%
42	Publication	Soleimani, N.. "Dietary supplementation of fructooligosaccharide (FOS) improves ...	<1%
43	Publication	Zhao, Juan, Yang Liu, Jun Jiang, Pei Wu, Weidan Jiang, Shuhong Li, Ling Tang, She...	<1%
44	Publication	"Posters", Clinical Microbiology and Infection, 4/2007	<1%
45	Publication	Haitham G. Abo-Al-Ela, Abeer F. El-Nahas, Shawky Mahmoud, Essam M. Ibrahim. "...	<1%

Immune response of saline tilapia (*Oreochromis niloticus*) to feed enriched with probiotic *Bacillus amyloliquefaciens* at high salinity

Rahmi R^{1*}, Fitri Indahyani², Akmal A³, Ifhan Dwinhoven⁴, Yuani Mundayana⁴, Muhammad Syaichudin³, Hidayat Suryanto S³, Nur Insana S¹, Khairun Nisaa³

¹Aquaculture Study Program, Faculty of Agriculture, Muhammadiyah University of Makassar, 90221, South Sulawesi, Indonesia.

²Aquaculture Study Program, Faculty of Agriculture, Muhammadiyah University of Pare pare, 91112, South Sulawesi, Indonesia.

³Research Center for Fisheries, National Research and Innovation Agency, Bogor Regency, 16915, West Java, Indonesia.

⁴Departement of Aquaculture, Fish Hatchery Technology Study Program, Pangkep State Polytechnic of Agriculture, 90655, South Sulawesi, Indonesia.

Abstract. Tilapia is a widely favoured fish due to its substantial meat content and appealing taste. The incorporation of probiotics in tilapia farming can enhance the fish's immunity against diseases. Additionally, probiotics can increase the total protein content of the feed and maintain the balance of digestive microbes. This study aimed to investigate the immune response of saline tilapia fed a diet enriched with the probiotic *B. amyloliquefaciens* at a salinity level of 10 ppt. The research focused on various blood parameters in tilapia, including total erythrocytes, leukocytes, and hematocrit levels, as well as the food conversion ratio (FCR) and specific growth rate (SGR). The rearing method involved maintaining approximately 100 g of tilapia fry for 45 days, with four treatments and three replications. The treatments included a control without *B. amyloliquefaciens* and additions of the probiotic at concentrations of 10^3 , 10^5 , 10^6 , and 10^7 CFU/mL. The study results indicated no statistically significant differences ($P>0.05$) in FCR values (1.16–1.24) and SGR (2.03–2.24% BW/day). However, the addition of *B. amyloliquefaciens* at a concentration of 10^7 CFU/mL led to an increase in erythrocyte count and hematocrit levels during the study period, with no observed effect on leukocyte count in saline tilapia reared at high salinity.

1 Introduction

* Corresponding author: rahmiperikanan@unismuh.ac.id

21 Tilapia (*Oreochromis niloticus*) is one of the most extensively cultivated fish species in Indonesia [1, 2]. In response to the growing demand for tilapia, farmers are exploring ways to enhance the health and growth of this fish in environments that may not be optimal, such as high-salinity conditions [3, 4]. Probiotics in fish are defined as living microorganisms that provide health benefits to their hosts, representing a promising solution [5, 6].

23 Excessive salinity levels can induce osmotic stress in fish, disrupting their internal homeostasis [7, 8]. This stress compromises the immune system, making fish more susceptible to diseases [9]. Research has shown that tilapia in high-salinity environments exhibit reduced activity of immune cells, such as lymphocytes and macrophages, which are essential for defense mechanisms.

10 Probiotics have been shown to enhance the immune response in fish through various mechanisms. They improve the health of the digestive tract, facilitating better nutrient absorption and overall health [10]. Additionally, probiotics contribute to the formation of beneficial microbiota in the intestines, which can inhibit pathogen growth [11]. Numerous studies have also demonstrated that probiotics can stimulate antibody production and enhance immune cell activity [12]. Probiotics offer an environmentally friendly alternative to antibiotics in aquaculture [13]. Erma's study [14] found that tilapia fed with probiotics under high salinity conditions had elevated levels of immunoglobulin M (IgM), a key antibody in the immune response. Moreover, fish fed with probiotics showed increased activity of T cells and B cells, which are vital for infection defense [15, 16].

36 Several studies have demonstrated that the administration of probiotics consisting of *Lactobacillus plantarum* to tilapia fish reared in high-salinity environments can result in an enhancement of immunoglobulin levels and lysozyme enzyme activity [17-19]. This evidence indicates that probiotics can enhance the immune system of fish, thereby increasing their resilience to environmental stressors, such as high salinity. This article will examine the impact of probiotic feed on the immune response of tilapia fish inhabiting high-salinity environments.

19 2 Materials and Methods

2.1 Preparation of fish

27 Prior to the commencement of the study, the tilapia seeds were acclimatised to a salinity of 15 ppt for a period of two weeks. The study employed a total of 12 containers, each with a capacity of 50 L and equipped with an aeration system. The stocking density of tilapia used was 20 fish with a length ranging from 7 to 9 cm, as reported by Rahmi et al. [20]. The fish were maintained in the containers for a period of 40 days, with feeding occurring three times a day (07.00, 12.00, 17.00) at a satiation level. Each container was filled with 20 liters of water at a salinity of 15 ppt and aerated to increase the dissolved oxygen levels within the maintenance medium.

2.2 Preparation of feed

The feed utilized in this research was formulated from a standard base and enhanced with *Bacillus amyloliquefaciens*, which was cultured in nutrient broth (NB). Initially, bacterial isolates were added to NB and incubated with agitation for 24 hours [21]. The bacterial suspension was then centrifuged at 6000 rpm for 10 minutes, after which the supernatant was discarded, and the bacterial pellet was resuspended in a saline solution. Subsequently, 1% of the feed was sprayed with this bacterial suspension [22, 23], air-dried, and then used.

2.3 Experimental Design

The experimental design employed was a completely randomized design (CRD) consisting of four treatments, each replicated three times. The treatments involved the addition of *Bacillus amyloliquefaciens* at concentrations of 10^3 CFU/mL (A), 10^5 CFU/mL (B), 10^7 CFU/mL (C), and 10^9 CFU/mL (D).

2.4 Fish performance indices

2.4.1 Blood profile

The total erythrocyte count was determined using the method described by Blaxhall and Daisley [24]. Blood samples, up to 0.2 mL, were collected using a red Sahli pipette marked to 0.5, followed by the addition of Hayem's solution up to the 101 mark. Blood from one to two fish was drawn using a 1 mL syringe. The blood in the pipette was then mixed thoroughly for 3-5 minutes, with 2-3 drops discarded to remove any unmixed blood. The sample was then placed onto a haemocytometer covered with a cover glass. The haemocytometer was examined under a microscope, and erythrocytes were counted in five visible sample boxes. The total erythrocyte count was calculated using the following formula:

$$\text{Total erythrocytes} = \text{number of erythrocytes counted} \times 10^4 \frac{\text{cells}^3}{\text{mm}} \quad (1)$$

The leukocyte count was conducted using the protocol outlined by Blaxhall and Daisley [24]. Blood samples mixed with EDTA in a vacutainer were drawn up to the 0.5 mark using a pipette, which was then cleaned with a tissue. The Turk solution was subsequently aspirated into the same pipette until it reached the 11 marks. The mixture was thoroughly homogenized for approximately three minutes to ensure complete dissolution. Before introducing the solution into the counting chamber, two or three drops were discarded. The solution was allowed to settle for one minute, after which leukocytes were counted using a microscope with $10\times$ or $40\times$ objective lenses. The leukocyte count was calculated using the following formula:

$$\text{Total leukocyte} = \text{number of erythrocytes counted} \times 10^4 \frac{\text{cells}^3}{\text{mm}} \quad (2)$$

Hematocrit levels were determined using the method described by Anderson and Siwicki [25]. Blood was drawn to fill $\frac{3}{4}$ of the microhematocrit tube. The end of the tube containing the blood was sealed with crytoceal by dipping it approximately 1 mm deep

to form a plug. The microhematocrit tube was then centrifuged at 5,000 rpm for 5 minutes. The length of the sedimented blood (a) and the total blood volume in the tube (b) were measured using a ruler. Hematocrit levels represent the proportion of blood cells (solids or sediment) in the blood fluid. The hematocrit value can be calculated using equation 3:

Haematocrit = (a/b) × 100% (3)

2.4.2 Growth performance and feed utilization

At the beginning and the end of the feeding trial, fish were weighed. The following fish performance indices were calculated:

Survival(%) = $\frac{\text{total of fish alive}}{\text{total of fish used}} \times 100$ (4)

Weight Gain (gr) = final weight – initial weight [26] (5)

Specific Growth Rate (SGR) = $100 \times \left[\frac{\ln(\sum \text{weight at the end of the study}) - \ln(\sum \text{weight at the beginning study})}{\text{rearing period (in days)}} \right]$ [27] (6)

Food Conversion Rate (FCR) = $\frac{\sum \text{weight at the end of the study} - \sum \text{weight at the beginning study}}{\text{Amount of feed consumed during the study period}}$ [28] (7)

3 Results

3.1 Blood Profile

The blood profile of saline tilapia fish in each treatment for each week of maintenance, as indicated by total erythrocytes (ranging from 9×10^5 to 3.3×10^6 cells/mm³) (Fig. 1), total leukocytes ($1,050 \times 10^3$ to $1,600 \times 10^3$ cells /mm³) (Fig. 2) and haemotocrit percentage (12.50 to 29.50%) (Fig. 3)




Fig. 1. Total of Erythrocytes observed over 35-day period in *Oreochromis niloticus*




Fig. 2. Total of Leukocytes observed over 35-day period in *Oreochromis niloticus*




Fig. 3. Percentage of Haematocrit observed over 35-day period in *Oreochromis niloticus*

3.2 Tilapia Growth and Feed Utilization Analysis

The survival rate of saline-treated tilapia ranged from 83.33% to 86.67%, with no significant differences observed ($P>0.05$). The feed conversion ratio (FCR) varied between 0.91 and 1.15 (Table 2). In contrast, the addition of *B. amyloliquefaciens* at a concentration of 10^5 CFU/mL (B) resulted in a significant increase ($P>0.05$) in body weight (BW), which ranged from 2.24 to 3.63 grams, and in specific growth rate (SGR), which ranged from 0.61% to 1.32% per day (Table 1).

Table 1. Survival, Weight Gain (WG), Specific Growth Rate (SGR), and Food Conversion Rate during observation in *Oreochromis niloticus*

Treatments	Survival (%)	WG (gr)	SGR(%/hari)	FCR
A	80.00±3.75 ^a	2.24±1.67 ^a	0.61±0.001 ^a	0.91±0.02 ^a
B	86.67±2.46 ^a	3.63±1.03 ^b	1.32±0.000 ^b	1.05±0.13 ^a
C	85.00±4.03 ^a	2.80±2.05 ^a	0.80±0.001 ^a	1.15±1.29 ^a

D	83.33±2.76 ^a	2.93±1.33 ^a	1.01±0.001 ^a	1.13±0.32 ^a
---	-------------------------	------------------------	-------------------------	------------------------

Note: Different superscripts in the same parameter indicate significantly different treatment effects (Tukey's test; P<0.05). Mean ± standar deviasi (SD), n=3

4 Discussion

The results of the study showed that there was a greater increase in the number of erythrocytes in the diet supplemented with *B. amyloliquefaciens* 10⁵ CFU/mL than in the other treatments (Fig. 1). The number of erythrocytes in this treatment increased significantly from day 7 to day 21, with the number of erythrocytes being 1.7×10⁶ cells/mm³ on day 7 and 3.3×10⁶ cells/mm³ on day 21. The erythrocyte count in this study was classified as normal condition. This is in line with the opinion of [28]; [29] that the normal range of total erythrocytes in tilapia is 2×10⁴-3×10⁶ cells/mm³. Changes in temperature and salinity also affect the number of erythrocytes in tilapia [30]. Changes in temperature and salinity can increase the number of erythrocytes if given continuously. This is because as temperature and salinity increase, the activity of oxygen uptake by erythrocytes increases and the fish's body will compensate for the changes in oxygen deprivation by increasing the number of erythrocytes.

The results of the study showed that the treatment with the addition of *B. amyloliquefaciens* probiotics for the treatment of 10⁵CFU/mL in saline tilapia feed did not show an increase in the number of leukocytes when compared to the treatment of 10³ CFU/mL, 10⁷ CFU/mL, and 10⁹ CFU/mL. It can be observed that the addition of *B. amyloliquefaciens* probiotics at a concentration of 10⁵ CFU/mL resulted in a non-significant increase in total leukocytes for each treatment over the course of the study period. This means that the addition of *B. amyloliquefaciens* probiotics can make the total leukocyte level from the beginning of the maintenance to the end of the maintenance of saline tilapia.

Leukocytes are integral to the immune response. In this study, leukocytes demonstrated no significant effect, despite being the most active component of the body's defense system, circulating in various forms within the bloodstream. Their primary function involves the destruction of infectious and toxic agents through phagocytosis and antibody formation [31]. The susceptibility of tilapia to disease at low temperatures can be attributed to fluctuations in leukocyte counts, either below or above normal levels. A reduced metabolic rate in fish leads to diminished immunity, increasing their vulnerability to diseases [32]. Elevated water temperatures enhance the immune response in fish. Variations in water quality during the growth period of tilapia, coupled with inadequate nutritional intake, can also result in decreased leukocyte counts, thereby compromising the immune system and increasing disease susceptibility [33, 34]. Additionally, increased temperatures can influence leukocyte counts in tilapia, leading to an elevation as part of the immune response to microorganisms.

The hematocrit levels observed during the study indicated that the treatment involved the addition of *B. amyloliquefaciens* probiotics at a concentration of 10⁵ CFU/mL exhibited a range of 15.45-33.70%. This value indicates the optimal result after 35 days

of study. The parameter that affects the measurement of red blood cell volume is the hematocrit. Haematocrit is defined as the percentage of red blood cell volume in the blood. Normal hematocrit values in tilapia range from 20 to 30%, with some marine fish exhibiting values of approximately 42% [35]. The hematocrit levels in the treatment based on the lowest value will gradually increase as the fish begin to adapt to the new environment and begin to obtain nutritional supplies from the food they eat. This is emphasized by [36], who states that hematocrit levels vary according to several factors, including nutritional intake, age, gender, body size and spawning period.

This suggests that the treatment, which includes the addition of probiotic *B. amyloliquefaciens* at a concentration of 10^5 colony-forming units (CFU) per milliliter, plays a pivotal role in enhancing the blood profile and overall resilience of tilapia. This is because it can boost the immune system response in tilapia through alterations in blood parameters, including total erythrocytes, total leukocytes, and haematocrit percentage, as observed during the course of the study. Furthermore, red blood cells are involved in the fish's innate and acquired immune response mechanisms, which contribute to the fish's body defense process. Erythrocytes can react to infections or other abnormalities within the fish's body [37, 38]. The hematocrit value in artificial feed supplemented with probiotics indicates that the synbiotic feed composition is optimal for the growth of tilapia, as it can elevate hemoglobin levels in the blood to attain normal values [23].

The survival rate of the saline tilapia (*O. niloticus*) was monitored throughout the duration of the study. A number of fish died in each treatment group. The variance analysis revealed that the inclusion of probiotics at varying doses in each treatment has no significant impact on the survival rate of saline tilapia (*O. niloticus*) ($P < 0.05$). However, Tukey's multiple comparison test indicated that the administration of different doses within the treatment had a significantly effect on the survival rate of saline tilapia.

The treatment involving the addition of probiotics, specifically *B. amyloliquefaciens* at a concentration of 10^5 CFU/mL, resulted in the highest body weight among saline tilapia. This outcome is hypothesized to be due to the probiotics meeting the necessary standards for effective feed digestion in saline tilapia. Previous studies have shown that incorporating probiotics into fish feed can increase body weight. This effect is attributed to the bacteria in probiotics, which improve digestion efficiency and nutrient absorption, thereby facilitating growth in both weight and length. Probiotics in feed can also enhance the nutritional profile, providing a source of easily digestible protein, biomass, and energy [39, 40, 41]. Additionally, probiotics can support production by promoting the growth and survival of cultured organisms [42].

The specific growth rate (SGR) was determined during the study, with the highest average treatment resulting from the addition of *B. amyloliquefaciens* probiotics at a density of 10^5 CFU/mL, exhibiting a rate of 1.32% per day. The analysis of variance (ANOVA) demonstrating that the administration of *B. amyloliquefaciens* probiotics had a statistically significant impact ($P < 0.05$) on the growth of saline tilapia (*O. niloticus*). The results of the Tukey post-hoc test demonstrated that there was a statistically significant difference between the treatment group that received the addition of *B. amyloliquefaciens* probiotics at a concentration of 10^5 CFU/mL and the treatment group that received the addition of *B. amyloliquefaciens* probiotics at other concentrations. The survival rate in saline tilapia demonstrated that the incorporation of *B. amyloliquefaciens*

probiotics into commercial feed resulted in a superior outcome compared to the use of feed lacking the probiotic supplementation. It is hypothesised that the addition of probiotics enhances immunity and improves the survival of fish. The study demonstrated that the number of leukocytes in tilapia receiving probiotics in their feed increased. These findings suggest that the administration of prebiotics may increase fish resistance to infection and stress, which could contribute to improved fish survival. A high number of leukocytes indicates a stronger immune response to various disorders. In a similar vein, [23] posited that the use of probiotics can increase the survival rate and resistance of fish to pathogen infections. This is corroborated by the findings of [43], who observed that treatment without the addition of probiotics resulted in a lower growth rate value compared to the treatment with the addition of probiotics.

The present study considers water quality parameters as one of the variables under observation. It is important to note that the conditions of normal water quality cannot be considered in isolation from the management of cultivation. Stable water quality conditions in several parameters, including temperature, pH, ammonia and salinity, have been demonstrated to correlate with optimal growth rates [51].

Table 2. Water quality parameters during period of study

Paramaters	Range of value	Optimum value	References
Temperature (°C)	28-30	25-30	[45, 46]
Salinity (ppt)	10-10,5	20	[47]
pH	7.10-8.00	6.5 – 8.5	[48, 49]
Ammonia (ppm)	0.0558-0.0776	0,1	[50]

In this study, water quality measurements were conducted over a 40-day period, encompassing temperature, pH, and salinity. Ammonia levels remained below 0.1 ppm throughout the maintenance phase, indicating a satisfactory status. These findings demonstrate that the incorporation of probiotic feed is an effective method for maintaining water quality at a consistent and optimal level, making it a suitable approach for the cultivation of saline tilapia.

5 Conclusion

The research results clearly show that adding probiotic *B. amyloliquefaciens* to saline tilapia feed at a concentration of 10⁵ CFU/mL optimises the total erythrocyte value, total leukocyte and haematocrit percentage. This is a significant finding that affects other treatments, including body weight and specific growth rate. The survival rate and feed conversion ratio of saline tilapia (*O. niloticus*) had no significant effect on saline tilapia during maintenance.

Acknowledgments

The authors would like to express their gratitude to the Ministry of Education, Culture, Research and Technology, Directorate General of Higher Education, Research and Technology for providing funding for the 2023 Research and Community Service Program (No. 0557/E5.5/AL.04/2023), as

well as to LP3M Muhammadiyah University of Makassar, Muhammadiyah University of Pare Pare, Hasanuddin University, and fellow students for their cooperation in implementing this research.

References

1. A.S. Abadi, I.B. Hismayasari, I. Supriatna, A. Yani, M. Sayuti, The mass death of tilapia (*Oreochromis niloticus*) in Sorong District, West Papua, Indonesia, Aq. Aqua. Cons. and Leg. **13**, 4, 1906-1916 (2020). <https://doi.org/10.47748/aac1.2020.1906-1916>
2. L. Emmy, Russ, Socio-Economic Development in Agriculture through Sustainable Practices, J. of Agr. and Soc-Eco. Sci. **117**, 9 (2021). <https://doi.org/10.18551/rjoas.2021-09>
3. A.A. El-Leithy, S.A. Hemeda, W.S.A. El Naby, A.F. El Nahas, S.A. Hassan, S.T. Awad, Z.A. Helmy, Optimum salinity for Nile tilapia (*Oreochromis niloticus*) growth and mRNA transcripts of ion-regulation, inflammatory, stress- and immune-related genes, Fish Physiol. and Biochem. **45**, 1217-1232 (2019). <https://doi.org/10.1007/s10695-019-00646-3>
4. M.E. Abd El-Hack, M.T. El-Saadony, M.M. Nader, H.M. Salem, A.M. El-Tahan, S.M. Soliman, A.F. Khafaga, Effect of environmental factors on growth performance of Nile tilapia (*Oreochromis niloticus*). International Journal of Biometeorology. **66**, 11, 2183-2194 (2022). <https://doi.org/10.1007/s00484-022-02347-6>
5. C. Sayes, Y. Leyton, C. Riquelme, Probiotic Bacteria as a Healthy Alternative for Fish Aquaculture, in: Antibiotic Use in Animals, InTech Publishers. (2018). <https://doi.org/10.5772/intechopen.71206>
6. S. Mondal, D. Mondal, T. Monda, J. Malik, Application of probiotic bacteria for the management of fish health in aquaculture. In Bacterial Fish Diseases, pp. 351-378, (Academic Press, 2022)
7. K. Zhou, Y. Huang, Z. Chen, X. Du, J. Qin, L. Wen, Y. Lin, Liver and spleen transcriptome reveals that *Oreochromis aureus* under long-term salinity stress may cause excessive energy consumption and immune response. Fish & Shellfish Immunology. **107**, 469-479 (2020). <https://doi.org/10.1016/j.fsi.2020.10.030>
8. N.A. Mohamed, M.F. Saad, M. Shukry, A.M. El-Keredy, O. Nasif, H. Van Doan, M.A. Dawood, Physiological and ion changes of Nile tilapia (*Oreochromis niloticus*) under the effect of salinity stress. Aquaculture Reports. **19**, 100567 (2021). <https://doi.org/10.1016/j.aqrep.2020.100567>
9. M. Abdel-Tawwab, M.N. Monier, S.H. Hoseinifar, C. Faggio, Fish response to hypoxia stress: growth, physiological, and immunological biomarkers. Fish Physiology and Biochemistry. **45**, 997-1013 (2019). <https://doi.org/10.1007/s10695-019-00614-9>
10. M.F. Rohani, S.M. Islam, M.K. Hossain, Z. Ferdous, Z. M.A. Siddik, M. Nuruzzaman, M. Shahjahan, Probiotics, prebiotics and synbiotics improved the functionality of aquafeed: upgrading growth, reproduction, immunity and disease resistance in fish. Fish & Shellfish Immunology. **120**, 569-589 (2022). <https://doi.org/10.1016/j.fsi.2021.07.011>

11. M.A. Dawood, S. Koshio, M.M. Abdel-Daim, H.V. Doan, Probiotic application for sustainable aquaculture. *Reviews in Aquaculture*. **11**(3), 907-924 (2019). <https://doi.org/10.1111/raq.12299>
12. F.K. Kuebutornye, E.D. Abarike, Y. Lu, V. Hlordzi, M.E. Sakyi, G. Afriyie, C.X. Xie, Mechanisms and the role of probiotic *Bacillus* in mitigating fish pathogens in aquaculture. *Fish Physiology and Biochemistry*. **46**, 819-841 (2020).. <https://doi.org/10.1007/s10695-019-00754-y>
13. W.A. Pereira, C.M.N. Mendonça, A.V. Urquiza, V.P. Marteinsson, J.G. LeBlanc, P.D. Cotter, R.P. Oliveira, Use of probiotic bacteria and bacteriocins as an alternative to antibiotics in aquaculture. *Microorganisms*. **10**(9), 1705 (2022). <https://doi.org/10.3390/microorganisms10091705>
14. K.W. Erma, Imunogenisitas dan tingkat keamanan vaksin polivalen streptococcus pada ikan nila *Oreochromis niloticus* (Linnaeus, 1758), (2023) <http://digilib.unila.ac.id/75896/>
15. E.B. Abarike, J. Cai, Y. Lu, H. Yu, L. Chen, J. Jian, F.K. Kuebutornye, Effects of a commercial probiotic BS containing *Bacillus subtilis* and *Bacillus licheniformis* on growth, immune response, and disease resistance in Nile tilapia, *Oreochromis niloticus*. *Fish & Shellfish Immunology*. **82**, 229-238 (2018). <https://doi.org/10.1016/j.fsi.2018.08.002>
16. A.M. Lusiastuti, S.D. Maryanti, U.P. Purwaningsih, Probiotik *Bacillus cereus* Untuk Pengendalian Penyakit Streptococcosis Pada Ikan Nila, *Oreochromis niloticus*. *Jurnal Riset Akuakultur*. **8**, 1, 109-119 (2014). <http://dx.doi.org/10.15578/jra.8.1.2013.109-119>
17. M.E. Abou-El-Atta, M. Abdel-Tawwab, N. Abdel-Razek, T.M. Abdelhakim, Effects of dietary probiotic *Lactobacillus plantarum* and whey protein concentrate on the productive parameters, immunity response, and susceptibility of Nile tilapia, *Oreochromis niloticus* (L.), to *Aeromonas sobria* infection. *Aquaculture Nutrition*. **25**(6), 1367-1377 (2019). <https://doi.org/10.1111/anu.12950>
18. G. Mohammadi, G. Rafiee, H.A. Abdelrahman, Effects of dietary *Lactobacillus plantarum* (KC426951) in biofloc and stagnant-renewal culture systems on growth performance, mucosal parameters, and serum innate responses of Nile tilapia *Oreochromis niloticus*. *Fish Physiology and Biochemistry*. **46**, 1167-1181 (2020). <https://doi.org/10.1007/s10695-020-00790-4>
19. M.S. Gewaily, M. Shukry, M.F. Abdel-Kader, M. Alkafafy, F.A. Farrag, E.M. Moustafa, M.A. Dawood, Dietary *Lactobacillus plantarum* relieves Nile tilapia (*Oreochromis niloticus*) juvenile from oxidative stress, immunosuppression, and inflammation induced by deltamethrin and *Aeromonas hydrophila*. *Frontiers in Marine Science*. **8**, 621558 (2021). <https://doi.org/10.3389/fmars.2021.621558>
20. Rahmi, Akmal, N.I. Salam, N, Optimasi ketahanan benih ikan nila salin (*Oreochromis niloticus*) terhadap infeksi streptococosis optimization of the resistance of tilapia salin (*Oreochromis niloticus*) seed against streptococosis infections. *Jurnal Galung Tropika*. **10**, 1 (2021). <https://doi.org/10.31850/jgt.v10i1.764>
21. I. Dwinhoven, T. Amalia, S. Haerunnisa, K.A. Bakar, Bioactive Compounds in Nest-Associated Bacteria Termites and Their Bioactivity as Antivibrio in Aquaculture

- Industry. International Journal of Applied Biology. **7**(2), 65-70. (2023).
<https://doi.org/10.20956/ijab.v7i2.31045>
22. Kurniawan, A., Suminto, S., Haditomo, A, Pengaruh Penambahan Bakteri Kandidat Probiotik *Bacillus methylotrophicus* pada Pakan Buatan terhadap Profil Darah dan Performa Pertumbuhan Ikan Nila (*Oreochromis niloticus*) yang Diuji Tantang dengan Bakteri *Aeromonas hydrophila*. Sains Akuakultur Tropis: Indonesian Journal of Tropical Aquaculture. **3**, 1, 82-92 (2019). <https://doi.org/10.14710/sat.v3i1.3956>
 23. Rahmi, R, Description of blood tilapia (*Oreochromis niloticus*) at high salinity fed with synbiotic feed. Jurnal Airaha. **12**, 2, 248-258. (2023).
<https://doi.org/10.15578/ja.v12i02.565>
 24. P.C. Blaxhall, K.W. Daisley, Routine Haematological Methods for Use with Fish Blood. Journal of Fish Biology. **5**, 3, 577–581 (1973). <https://doi.org/10.1111/j.1095-8649.1973.tb04510.x>
 25. D.P. Anderson, A.K. Siwicki, Basic hematology and serology for fish health programs. Paper presented in the second symposium on disease in Asian Aquaculture Aquatic Animal Health and Environment. **1**, 1 (1993)
 26. M.I. Effendie, Metode Biologi Perikanan, (Yayasan Dewi Sri Bogor, 1979)
 27. E.A. Huisman, Principles of fish production. Department of Fish Culture and Fisheries, (Wageningen Agriculture University. Wageningen. Netherland, 1987)
 28. R. Hartika, M. Mustahal, A.N. Putra, Gambaran darah ikan nila (*Oreochromis niloticus*) dengan penambahan dosis prebiotik yang berbeda dalam pakan. J. Perikanan dan Kelautan. **4**, 4 (2014). <http://dx.doi.org/10.33512/jpk.v4i4.174>
 29. N. Subryana, Wardiyanto, O. Susanti, Penggunaan ekstrak daun kelor *Moringa oleifera* (Lam, 1785) untuk meningkatkan imunitas non spesifik benih ikan nila *Oreochromis niloticus* (Linnaeus, 1758) yang diinfeksi *Aeromonas hydrophila*. J. of Aqua. and Fish Health. **9**, 3 (2020). <https://doi.org/10.20473/jafh.v9i3.16321>
 30. Y. Cahyanti, I. Awalina, Studi literatur: pengaruh suhu terhadap ikan nila (*Oreochromis niloticus*). Panthera: J. Il. Pend. Sains dan Terapan. **2**, 4 (2022).
<https://doi.org/10.36312/pjipst.v2i4.110>
 31. P.R. Machado, M.I.A. Araújo, L. Carvalho, E.M. Carvalho, Immune response mechanisms to infections. Anais Brasileiros de Dermatologia. **79** (2004).
<https://doi.org/10.1590/S0365-05962004000600002>
 32. C. Pohlenz, D.M. Gatlin III, Interrelationships between fish nutrition and health. Aquaculture. **431** (2014). <https://doi.org/10.1016/j.aquaculture.2014.02.008>
 33. J. Raa, The use of immune-stimulants in fish and shellfish feeds. In: Cruz-Suarez, L. E., Ricque-Marie, D., Tapia-Salazar, M., Olvera-Novoa, M. A., Civera-Cerecedo, R. (Eds.), *Avances en Nutrición Acuicola V*, Memorias del V Simposium Internacional de Nutrición Acuicola, Mérida, Yucatan, México. pp. 47–56 (2000)
 34. Kiron, V. Ani, Fish immune system and its nutritional modulation for preventive health care. Feed Sci. and Tech. **173**, 2 (2012).
<https://doi.org/10.1016/j.anifeeds.2011.12.015>
 35. C.E. Bond, 1979. Philadelphia: Saunders College Publishing, 514p.

36. Y. Kuswardani, Pengaruh pemberian resin lebah terhadap gambaran darah maskoki *Carassius auratus* yang terinfeksi bakteri *Aeromonas hydrophila*. Master Thesis, Bogor Agricultural Institut, 2006
37. M. Witeska, Erythrocytes in teleost fishes: a review. *Zoology and Ecology*. **23**, 4 (2013). <https://doi.org/10.1080/21658005.2013.846963>
38. K.R. Grant, Fish hematology and associated disorders. *Vet. Clinics: Exotic Animal Practice*. **18**, 1 (2015). <https://doi.org/10.1016/j.cvex.2014.09.007>
39. S. Aslamyah, Kualitas Lingkungan dan Aktivitas Enzim Pencernaan Udang vannamei (*Litopenaeus vannamei*) pada Berbagai Konsentrasi Probiotik Bioremediasi *Bacillus* sp. *Fish Scientiae*. **1**, 2 (2011). <https://doi.org/10.20527/fishscientiae.v1i2.19>
40. J.E. Setiawati, Y.T. Tarsim, Adiputra, H. J. Siti, Pengaruh Penambahan Probiotik pada Pakan dengan Dosis Berbeda terhadap Pertumbuhan, Kelulushidupan, Efisiensi Pakan dan Retensi Protein Ikan Patin (*Pangasius hypophthalmus*). *Rekayasa dan Teknologi Perairan*. **1**, 2 (2013).
41. J.F. Sumule, T.T. Desiana, Rusaini, Aplikasi Probiotik pada Media Pemeliharaan terhadap Pertumbuhan dan Sintasan Ikan Nila Merah (*Oreochromis* sp.). *J. Agrisains*. **18**, 1 (2017)
42. T.W. Hapsari, M.A. Tjahjaningsih, Alamsjah, H. Pramono, Aktivitas Enzimatis Bakteri Proteolitik Asal Gastrointestinal Udang Vannamei (*Litopenaeus vannamei*). *J. of Mar. and Coastal Sci*. **5**, 3 (2000)
43. A.H. Primashita, B.S. Rahardja, Prayogo, Pengaruh Pemberian Probiotik Berbeda Dalam Sistem Akuaponik Terhadap Laju Pertumbuhan Dan Survival Rate Ikan Lele (*Clarias* Sp.). *J. of Aqua. Sci*. **1**, 1 (2017)
44. M. Salsabila, H. Suprpto, H. Teknik pembesaran ikan nila (*Oreochromis niloticus*) di instalasi budidaya air tawar pandaan, Jawa Timur *J. of Aqua. and Fish Health*. **7**, 3 (2018)
45. H. Ariadi, B.D. Madusari, D. Mardhiyana, D. Analisis pengaruh daya dukung lingkungan budidaya terhadap laju pertumbuhan udang vaname (*L. vannamei*). *Enviro. Scienteae*. **18**, 1 (2022). <http://dx.doi.org/10.20527/es.v18i1.12976>
46. A.N. Bergman, Physiology of the Lake Magadi tilapia, *Alcolapia grahami*, a fish living in an extreme environment of high pH, temperature and alkalinity and frequent hypoxia, Ph.D. thesis, Univ. of Wyoming, Department of Zoologi and Physiology (2000)
47. N. Montoya-Camacho, F.J. Castillo-Yáñez, E. Márquez-Ríos, S. Ruíz-Cruz, A.A. Arvizu Flores, H.J. Barrales-Cureño, V.M. Ocaño-Higuera, Evaluation of biochemical, chemical, physical and microbiological quality of tilapia (*Oreochromis niloticus*) muscle during 0 and 5° C storage. *Biotecnia*. **23**, 2 (2021). <https://doi.org/10.18633/biotecnia.v23i2.1418>
48. A.A.A. El-Leithy, S.A. Hemeda, W.S.A El Naby, A.F. El Nahas, S.A. Hassan, S.T. Awad, Z.A. Helmy, Optimum salinity for Nile tilapia (*Oreochromis niloticus*) growth and mRNA transcripts of ion-regulation, inflammatory, stress-and immune-related genes. *Fish Phys. and Biochem*. **45** (2019). <https://doi.org/10.1007/s10695-019-00640-7>

49. K. Zhou, T.T. Cui, P. L. Li, N.J. Liang, S.C. Liu, C.W. Ma, Z.H.J, Modelling and predicting the effect of temperature, water activity and pH on growth of *Streptococcus iniae* in Tilapia. Peng., J. of Applied Micro. **105**, 6 (2008).
<https://doi.org/10.1111/j.1365-2672.2008.03969.x>
50. N. Arpi, Fahrizal, M. Novita, Isolation of fish skin and bone gelatin from tilapia (*Oreochromis niloticus*): Response surface approach. In IOP Conference Series: Materials Science and Engineering. **334**, 1 012061 (2018).
<https://doi.org/10.1088/1757-899X/334/1/012061>
51. R.V. Constantino, R.M.C. Templonuevo, L.J. Fajardo, Stress responses of red tilapia (*Oreochromis spp.*) to high ammonia levels. International J. of Fish, Aqu. Stu. **7**, 6 (2019)